



Stability of Active Constituents of Hops (*Humulus lupulus*) Strobiles and their Ethanolic Extracts during Storage

**Daniel Gagnon^{1*}, Chitra Wendakoon², Robert Smith³
and Jeremy Leker³**

¹Herbs, Etc., Inc., Santa Fe, NM 87507, USA.

²Independent Consultant, Austin, TX, USA.

³S.S. Steiner, Yakima, WA, USA.

Authors' contributions

*This work was carried out in collaboration between all authors. Author DG designed the study, wrote the protocol, supervised the preparation of the *H. lupulus* samples and helped in revising the manuscript. Author CW provided advice throughout the project and prepared the final manuscript. Authors RS and JL performed the HPLC analysis of the *H. lupulus* samples. All authors read and approved the final manuscript.*

Original Research Article

Received 6th May 2014

Accepted 30th June 2014

Published 20th July 2014

ABSTRACT

Aims: The purpose of this study was to evaluate the stability of three major active constituents (humulones, lupulones and xanthohumol) in dried hops (*Humulus lupulus*) strobiles (whole and ground) as well as their ethanolic extracts during storage.

Methodology: A comparative study of humulones, lupulones and xanthohumol levels of *H. lupulus* strobiles during storage was carried out. Dried whole strobiles and cryogenically ground dried strobiles stored at -15°C as well as ethanol extracts of the strobiles prepared using different ethanol concentrations (10%, 30%, 50%, 70%, and 95%) and stored at room temperature, were analyzed by HPLC to quantify each constituent. These hops samples were analyzed immediately after preparation, and then one year and two years later to determine the concentrations of the constituents.

Results: HPLC analysis indicated that the amount of all three constituents in the ground strobiles and in the ethanol extracts decreased gradually during the storage period. The 10% and 30% ethanol extracts had very low amounts of constituents initially and were

*Corresponding author: Email: botandan@aol.com;

practically devoid of constituents at the end of two years. The 50% ethanol extract contained considerable amounts of humulones and xanthohumol, and low levels of lupulones initially, but lost substantial amounts over time. The 70% and 95% ethanol extracts showed higher levels of all three constituents, while the 95% *H. lupulus* ethanol extract contained the highest constituent levels throughout the experimental period. The ethanol content of the extract had a direct correlation to the constituent levels; the higher the ethanol level, the higher the initial and subsequent constituent levels.


Conclusion: Both dried hops and ethanol extracts lose active components over storage time. When preparing extracts, at least 70% ethanol is necessary to extract the highest levels of three bioactive constituents and to retain them over a two-year period. Ethanol concentration is a critical factor to be considered in hops extraction process.

Keywords: Hops; ethanol extracts; humulones; lupulones; xanthohumol; storage stability.

1. INTRODUCTION

Humulus lupulus L., commonly known as hops, belongs to the Hemp [*Cannabaceae*] family [1] and has been cultivated throughout the temperate regions of the world primarily for the brewing industry [2]. The strobiles (also known as cones) produced by female plants of hops are the desired parts for the brewing industry. For this reason, only the female plants are cultivated, and the strobiles are harvested during late summer for further processing [3]. The brewing industry generally grows *H. lupulus* from "root" cuttings and not from seeds since this cultivation method maintains a genetically consistent product. It also aids in controlling the aroma characteristics and the amount of active constituents found in the strobiles [4]. However, the overall strobile chemical composition still depends on a multitude of factors including variety, growing region, growing conditions, harvesting time, as well as drying and storage conditions [5,6].

Oleoresin glands present in the strobiles produce a resinous yellowish/reddish powder called lupulin [7]. Numerous compounds present in lupulin are of economical interest. The volatile oils and bitter acids are the most significant classes of compounds in terms of economic value. Two of the major constituents found in lupulin resin are humulones (alpha acids) and lupulones (beta acids) [8], which are phloroglucinol derivatives. Xanthohumol, another bioactive flavonoid, has also been isolated from the hops resin [9].



The stability of the constituents of the strobile pellets or extracts used in the brewing industry has been of great importance [10]. Moreover, the stability of humulones in *H. lupulus* during storage of strobiles has been recognized as a critical issue since humulones provide most of the bitterness in beer [11]. When the hop cones are harvested, the moisture content is around 75-80% and, in order to prevent deterioration, reduction of moisture content is necessary before storage or processing. Skinner et al. [12] demonstrated that the

constituents' rate of deterioration was related to the storage temperature; that every 15°C rise in the storage temperature doubled the deterioration rate. In order to prevent the loss of hops' active constituents, it is important to store the strobiles at a low temperature; preferably below 0°C. Weber et al. [13] studied the effects of post-harvest handling on the quality and storage stability of strobiles. Their study showed that decreasing the kilning temperature, using a lower compression force during bundling of strobiles, and wrapping the strobiles in burlap instead of plastic were instrumental in producing a superior product. These results concluded that, of the three factors tested (temperature, compression of the strobiles, and the material used to wrap the strobiles), elevated temperature had the most negative influence.

Clinical herbalists frequently use hops to treat a variety of ailments [14]. In England, *H. lupulus* strobiles have been recommended for their skin anti-infective properties for hundreds of years [15] and the strobiles have been used as a wash for impetigo, boils, and abscesses [16]. In traditional European folk medicine, *Humulus lupulus* was frequently mentioned as an infusion or a fomentation to treat skin sores, cuts, and injuries [17-19]. More recently, Bartram [20] suggested that the antimicrobial properties of hops could be helpful to treat skin infections. At present, however, *H. lupulus* is mainly used in modern phytotherapy for its nervous system sedative effects, to stimulate gastric secretions, and to improve digestive function [21]. Additionally, *H. lupulus*' phyto-estrogenic properties [22] and antiviral activities [23] have recently been investigated and ascribed to its constituents.

It is generally acknowledged that clinical results can only be achieved if the herbs' active constituents are present in sufficient quantity to reach therapeutic levels [24]. This important issue, although of considerable concern to practitioners, has not been properly addressed by researchers [25]. Additionally, research suggests that the ethanol percentage used to extract herbs has a significant impact on the amount of active constituents found in the final extract [26]. In some instances, a high percentage of ethanol yields higher levels of active constituents [27] while, in other instances, a low ethanol level actually yields higher levels of active constituent [28]. The present study focuses on the amounts of bioactive constituents present in dried and ground *H. lupulus* strobiles and the extracts made with varying ethanol concentration and the storage stability of these constituents over time.

2. MATERIALS AND METHODS

2.1 Plant Materials

Humulus lupulus L. plant materials used in this study consisted of whole dried strobiles of the Super Galena variety grown in the Yakima Valley, Washington, USA (Hopsteiner, a division of S.S. Steiner, New York, NY). The strobiles were collected in the autumn of 2009, dried, and stored in warehouses under frozen conditions until shipped to Herbs, Etc., Inc. where they were stored in a freezer maintained at -15°C. Identity of the material was confirmed by the first author using macroscopic and organoleptic methods. A voucher specimen of whole strobiles was stored in a freezer maintained at -15°C.

2.2 Solvents

Five concentrations of ethanolic extracts (10%, 30%, 50%, 70%, and 95%) were prepared using 95% USP grade ethyl alcohol (Pharmco-Aaper, Shelbyville, KY) and water. The ethanol concentrations were verified using a hydrometer. A sample of each ethanol

concentration was set aside in an amber-colored glass bottle as reference material and stored at room temperature in a dark closet for solvent control studies.

2.3 Preparation of Hops Strobile Extracts

2.3.1 Cryogenic grinding

In order to prevent the loss of heat-sensitive constituents, the strobiles were cryogenically ground using a hammer mill (Fitzpatrick Manufacturing, Sterling Heights, MI) cooled by the injection of USP-grade liquid nitrogen into the grinding chamber. Samples of ground strobiles were then stored at -15°C.

2.3.2 Cold-process percolation extraction

On the same day that the *H. lupulus* strobiles were powdered using the above-described cryogenic grinding method, a cold-process percolation extraction method using a 1:5 herb-to-solvent (ethanol) ratio was used to extract the ground strobiles. Five different concentrations of ethanolic extracts (10%, 30%, 50%, 70%, and 95%) were used as solvents. The finished products were filtered to remove the sediments present in the liquid extracts and stored in amber-colored glass bottles. Three lots of samples were set aside, a) One lot for immediate analytical purposes, b) The second lot as reserve samples for quality assurance purposes and c) An additional lot for analytical purposes at the end of one year and two year storage. All the ethanol extract samples were kept at room temperature in a dark closet.

2.4 Chemical Analysis

Chemical analysis of the whole and ground strobiles stored at -15°C as well as the five ethanolic extracts prepared from the same strobile lot and stored at room temperature was performed by S.S. Steiner, in Yakima, WA. The amount of the three bioactive constituents [humulones, lupulones, and xanthohumol] in each sample was quantified using an HPLC method. The samples were analyzed again one year and two years later to determine the changes in constituents during storage.

2.4.1 Standards and sample preparation for HPLC analysis

Working standard solutions of humulones and lupulones were prepared by bath sonication of 0.5g (measured to 0.1mg) of the international calibration extract (ICE-3) obtained from American Society of Brewing Chemists (ASBC, St. Paul, MN). The ICE-3 extract was dissolved in 50 ml of methanol and diluted (1:10) in acidic methanol (0.5 ml of 85% phosphoric acid in 1 liter of methanol). The standard solution of xanthohumol was prepared by dissolving approximately 20mg of xanthohumol (in-house standard) in 100 ml of acidic methanol. Samples were diluted (1:20) with acidic methanol prior to analysis.

2.4.2 HPLC analysis

Quantification of the three constituents, humulones, lupulones, and xanthohumol, were carried out by the methods described by the European Brewing Congress, method EBC 7.7, and the American Society of Brewing Chemists, method ASBC HOPS-14, using a Shimadzu HPLC system equipped with diode array detector. The mobile phase was composed of

72.5% methanol, 26.5% water, 0.85% phosphoric acid, and 0.075mM Sodium EDTA. A C-18 column, Kinetex 2.6um, 4.6x50mm (Phenomenex) was used to separate the compounds. The flow rate was adjusted to 1.3 ml/minute at 40°C, and 10µl of samples and calibration solutions were injected into the column. The detector wavelength was set at the absorbance of 270 nm for humulones and lupulones, and at 367 nm for xanthohumol.

3. RESULTS AND DISCUSSION

3.1 HPLC Analysis of Whole and Ground *H. lupulus* Strobiles before and after Storage

3.1.1 Initial analysis

The HPLC results revealed that the whole dried *H. lupulus* strobiles initially contained 11.4g of humulones, 7.4g of lupulones, and 0.46g of xanthohumol per 100 grams of strobiles (Table 1). Immediately after the grinding process, the cryogenically-ground strobiles from the same lot were shown to contain lesser amounts of the three constituents. Even though the strobiles were ground using ultra-cold cryogenic technology, they were shown to have lost 8.8% of their humulones and 7.2% of their lupulones (Table 2). With a net 2.2% loss, xanthohumol showed the smallest constituents loss of all.

Fig. 1 shows the HPLC chromatogram of the standard compounds used in the analysis. The total humulones (cohumulone plus n+adhumulone), total lupulones (colupulone plus n+adlupulone) and xanthohumol levels were determined using the ICE-3 standard.




Fig. 1. HPLC chromatogram of ICE-3 standards showing humulones (alpha acids) lupulones (beta acids) and xanthohumol

One year later. All the three constituents were found to be decreased in both whole and ground strobiles stored at -15°C after one year (Table 1). The total amount of humulones found in whole strobiles was slightly higher than the amount found in ground strobiles (9.9g/100g vs. 9.7g/100g) after one year of storage, while the amount of lupulones was higher in ground strobiles than in whole strobiles (6g/100g vs. 5.6g/100g). The amount of

xanthohumol found in both whole and ground strobiles was exactly the same (0.43g/100g) after one year of storage (Table 1).

Table 1. HPLC analysis of dried whole and cryogenically-ground *Humulus lupulus* strobiles (stored at -15°C) immediately after grinding, one year, and two years after the grinding process

Constituents (%)	Storage time					
	Whole strobiles			Ground strobiles		
	Initial	Year 1	Year 2	Initial	Year 1	Year 2
Humulones	11.40	9.90	11.67	10.40	9.66	7.98
Lupulones	7.48	5.64	6.78	6.94	6.03	5.34
Xanthohumol	0.46	0.43	0.43	0.45	0.43	0.35

When comparing the percentage loss of whole and ground strobiles, the approximate amount of humulones (13% vs. 7%), lupulones (25% vs. 13%) and xanthohumol (7% vs. 5%) lost in the first year of storage was greater in the whole strobile sample than in the ground sample (Table 2).

Table 2. Percentage loss of constituents in whole and ground strobiles (stored at -15°C) immediately after grinding (initial), after one year and two years of storage

Constituents (%)	Percentage loss					
	Whole strobiles			Ground strobiles		
	Initial	Year 1	Year 2	Initial	Year 1	Year 2
Humulones	0	13.2	0	8.8	7.1	23.3
Lupulones	0	24.5	9.4	7.2	13.1	23.0
Xanthohumol	0	6.5	6.5	3.0	4.4	22.2

Two years later. After two years of storage at -15°C, whole hops strobiles were found to contain higher levels of humulones and lupulones, while in ground samples, the levels were decreased (Table 1). The anomaly of higher levels of constituents in whole hops after two years of storage may stem from the fact that each individual strobile is highly variable in its active constituent levels, while ground strobiles are a mixture of many ground and homogenized strobiles. Different strobiles used in the HPLC analysis may be the reason for this higher amount of humulones and lupulones. The ground strobiles showed an almost 23% decrease in humulones and lupulones as well as a 22% decrease in xanthohumol levels compared to the initial levels (Table 2). The ground strobiles powder tended to show more uniformity in their levels of active constituents.

3.2 HPLC analysis of Ethanol Extracts of *H. lupulus* Strobiles Immediately after Extraction

Fig. 2 shows the HPLC chromatogram of the 95% ethanol extract. All ethanol extracts were analyzed under the same conditions, and the amount of each constituent was calculated using the standards.




Fig. 2. HPLC chromatogram of the 95% ethanol extract

3.3 Initial Analysis

The results from the chemical analysis of *H. lupulus* ethanolic strobile extracts showed that humulones (2,120mg/100ml), lupulones (1,440mg/100ml) and xanthohumol (90mg/100ml) were highest in the 95% ethanolic extract (Table 3). There was a striking difference between the amount of active constituents reported in the 10% ethanolic extract and the 95% ethanolic extract. The 10% ethanolic extract had 53 times less alpha acids, 96 times less beta acids, and 45 times less xanthohumol than the 95% ethanolic extract.

Table 3. Changes in the amounts of bioactive constituents of *Humulus lupulus* strobiles ethanol extracts (stored at room temperature) immediately after extraction, one year, and two years after storage

Ethanol %	Constituents (mg/100ml)								
	Humulones			Lupulones			Xanthohumol		
	Initial	Year 1	Year 2	Initial	Year 1	Year 2	Initial	Year 1	Year 2
10	40	3	0.8	15	0.4	ND	2.0	0.09	ND
30	60	16	6.8	17	0.5	0.1	2.0	0.4	0.3
50	630	520	404	53	36	30	26	18	9.6
70	2,010	1,810	1,500	1,050	1,000	892	78	70	43
95	2,120	2,030	1,660	1,440	1,370	1,170	90	97	59

These results clearly demonstrate that, as the amount of ethanol in the menstruum used to extract the strobiles increased, the level of active constituents extracted also increased. The biggest increase in humulones level occurred when the ethanol percentage increased from 30% to 50%; the humulones amount increased from 60 to 630mg/100ml, a ten-fold increase. A similar increase in xanthohumol levels was observed when the ethanol percentage increased from 30% to 50%; the amount of xanthohumol increased from 2 to 26mg/100ml, a thirteen-fold increase. Conversely, the biggest increase in the percentage of lupulones occurred when the ethanol content increased from 50% to 70%; at that level, the amounts of lupulones increased from 53 to 1,050mg/100ml, a twenty-fold increase of lupulones. At an

ethanol concentration of 70%, or greater, a substantial increase in the amount of all three constituents present in the extract was noted (Table 3).

3.4 HPLC analysis of *Humulus lupulus* Ethanol Extracts after one Year and two Years of Storage

HPLC analysis of the five ethanol extracts stored at room temperature for 2 years showed that the different ethanol levels used to extract the strobiles greatly influenced the stability of each constituent over time (Table 3).

3.4.1 One year later

The 10% *H. lupulus* ethanol extract, which began with 40mg/100ml humulones was found to contain only 3mg/100ml of the constituent one year later indicating a loss of 92.5% of humulones. In comparison, not only did the 95% ethanol extract start with a much higher level of humulones (2,120mg/100ml), but one year later, the amount of humulones remained at 2,030mg/100ml, representing a relatively small loss of 4.25%.

Overall, lupulones levels showed a greater loss over a one year period when compared to humulones. The 10% *H. lupulus* ethanolic extract initially contained 15mg/100ml of lupulones but one year later, only 0.4mg/100ml of lupulones remained. This represents a loss of 97.5% lupulones in one year. Conversely, the 95% ethanolic extract started with a lupulones level of 1,440g/100ml and one year later was still found to contain 1,370mg/100ml of lupulones. The 95% ethanol extract had lost only 5% of its lupulones, while the 10% ethanol extract had lost most of its lupulones.

Xanthohumol analysis results showed a smaller loss when compared to humulones and lupulones. The 10% ethanol extract initially contained 2mg/100ml of xanthohumol. One year later, HPLC analysis revealed that it contained 0.09mg/100ml, a loss of nearly 96%. However, the 95% ethanol extract started with a xanthohumol level of 90mg/100ml and ended with a level of 81mg/100ml indicating a loss of only 10%.

Two years later. The 10% ethanol extract showed 0.8mg/100ml of humulones, while the 95% ethanol extract contained 1,660mg/100ml of humulones (Table 3). The 10% ethanol extract lost 98% of its humulones during the 24 month period, while the 95% ethanol extract lost only 22% of its humulones. The HPLC analysis revealed that, the lower the ethanol level used to make the *H. lupulus* ethanolic extract, the greater the loss of humulones over time. As the ethanol content used to extract the strobiles increased, higher levels of humulones were also retained over time. The higher ethanol content seemed to act as a preservative of humulones.

After a two year storage period, the lupulones were not detectable in the 10% ethanol extract, while the 95% ethanol extract still contained 1,170mg/ml or 81% of its lupulones. In effect, the 10% ethanol extract lost 100% of its lupulones, while the 95% ethanol extract lost only 19% of its lupulones. At the end of two years, analysis of xanthohumol levels revealed that there were no detectable levels of this constituent in the 10% ethanol extract, while the 95% ethanol extract had levels of 59mg/ml or 52% of the initial levels of xanthohumol.

When lower levels of ethanol were used to extract *H. lupulus* strobiles, concomitant lower levels of active constituents were extracted. Conversely, the higher the ethanol levels used

to extract the strobiles, the less active constituents were lost over a one year or two year storage period. A summary of the percentage of active constituents remaining in the ethanolic extracts over a one and two year period is shown in Table 4. The 95% *H. lupulus* ethanolic extract lost less than 5% of both humulones and lupulones over a one year period. However, the 10% ethanolic extract lost more than 92% and 97% respectively of the same two compounds. Over a two-year storage period, the 95% ethanolic extract lost 25% and 15% of its humulones and lupulones, while the 10% ethanolic extract lost over 99% and 100% of these two constituents.

When the stability of the three major active constituents is compared over time, it was observed that there was a strong correlation between the amount of ethanol used to extract the strobiles and the amount of active constituents retained in the extract after a storage period of one and two years. Similar results have been obtained in previous research for other herbs including hops [26]. One observation is that the higher the level of ethanol used to extract *H. lupulus* strobiles, the better the stability of the constituents in the ethanol extract stored at room temperature over time. The two extracts that contained the most constituents were the 70% and 95% ethanol extract; they both retained approximately three-quarter of their humulones and lupulones over a two year period. The extract that contained 50% ethanol lost substantial levels of constituents, while the two extracts with 10% and 30% ethanol lost virtually all of their constituents. These last two extracts are of particular concern in that they did not contain significant levels of constituents to start with, and then, proceeded to lose all of them over the two year storage period. The present study showed for the first time that ethanol concentration is a very important factor to be considered in hops extraction process and storage.

Table 4. Percentage of constituents remaining in *Humulus lupulus* ethanolic extracts after one year and two years of storage at room temperature (21°C)

Ethanol (%)	Constituents (%)					
	Humulones		Lupulones		Xanthohumol	
10	7.5	0.2	2.7	0	4.5	0
30	26.7	11.3	2.9	0.6	20	15
50	82.5	64.1	67.9	56.6	76	36.9
70	90.1	74.6	95.2	85.0	103	55.1
95	95.8	78.3	95.1	81.3	90	65.6

4. CONCLUSION

Whole dried *Humulus lupulus* strobiles contained the constituents humulones, lupulones and xanthohumol at 11.4%, 7.5%, and 4.6% levels respectively. After grinding the strobiles under cryogenic conditions, the levels of humulones and lupulones were reduced by nearly 10%. During the storage for two years at -15°C, the levels were further reduced. The *Humulus lupulus* strobiles extracted with an ethanol level of 70% or higher yields more active constituents than extracts made with lower concentrations of ethanol. Further, ethanol levels of 70% or higher helps to preserve the active constituents found in *H. lupulus* extracts for a longer period of time (at least over a two-year period) than extracts made with lower ethanol levels. The data obtained in this study may be of help to clinical herbalists as well as to the dietary supplements or pharmaceutical industry in developing nutraceutical and pharmaceutical products using *H. lupulus* extracts for human ailments. It is recommended that clinical herbalists use at least 70% *H. lupulus* ethanolic extracts in their clinical practice. Thus, they will gain an additional assurance that the *H. lupulus* ethanolic extract dispensed

to their clients contains the level of the constituents necessary to achieve the clinical results desired.

CONSENT

Not applicable.

ETHICAL APPROVAL

Not applicable.

ACKNOWLEDGEMENTS

The first author would like to thank the staff of Herbs, Etc. for grinding the *H. lupulus* strobiles and for manufacturing the various concentrations of ethanolic extracts from the ground strobiles.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. USDA, ARS, National Genetic Resources Program. Germplasm Resources Information Network - (GRIN) [Online Database]. National Germplasm Resources Laboratory, Beltsville, Maryland. Accessed 4 February 2013. Available at: http://www.ars-grin.gov/cgi-bin/npgs/html/tax_search.pl.
2. Biendl M, Pinzl C. Hops and Health: Uses, Effects, History, German Hop Museum, Wolnzach, Germany; 2008.
3. Koetter U, Biendl M. Hops (*Humulus lupulus*): A Review of its Historic and Medicinal Uses. Herbal Gram. 2010;87:44-57.
4. Hop Steiner Group. Guidelines for Hop Buying. New York, NY: SS Steiner, Inc; 2008.
5. Virant M, Majer D. Hops storage index – Indicator of a brewing quality. International Hop Growers Convention; 2006. Accessed 30 September 2012. Available at: <http://midlandscraftbrewing.aspx2.dotnetpanel.net/Technical%20information/Hop%20Storage%20Index.pdf>.
6. Forster A. The quality chain from hops to hop products. EBC Proceedings 29th Congress. 2003;17:156-165.
7. Evans W. Trease and Evans' Pharmacognosy. 13th edition. London, England: Bailliere Tindall; 1989.
8. Verzele M. Centenary review. 100 years of hop chemistry and its relevance to brewing. J. Inst. Brew. 1986;92:32-48.
9. Gerhauser C. Broad spectrum and anti-infective potential of xanthohumol from hop (*Humulus lupulus* L.) in comparison with activities of other hop constituents and xanthohumol metabolites. Molecular Nutrition and Food Research. 2005;49(9):827-831.
10. Garetz M. Hops storage: How to get – and keep – your hops' optimum value. Brewing Techniques. 1994;2(1):26-32.

11. Smith B. Brewing Hops Storage: Preserving Precious Hops. 2008. Accessed 10 December 2011. Available at: <http://www.beersmith.com/blog/2008/04/15/brewing-hops-storage-preserving-precious-hops/>.
12. Skinner RN, Hildebrand RP, Clarke BJ. The effect of storage temperature on the stability of the alpha-acids content of baled hops. J Inst Brew. 1977;83:290-294.
13. Weber KA, Janguaard NO, Foster RT. Effect of postharvest handling on quality and storage stability of Cascade hops. American Society of Brewing Chemists. 1979;37(2):58-60.
14. Hoffmann D. Medical Herbalism. Rochester, Vermont: Healing Arts Press; 2003.
15. Culpeper N. Culpeper's Complete Herbal. Champaign. IL: Book Jungle; 1999.
16. Grieve M. A Modern Herbal. New York, NY: Dover Publications, Inc; 1971.
17. Van Hellemont J. Compendium de Phytotherapie. Brussels, Belgium: Association Pharmaceutique Belge; 1986. French.
18. Valnet, J. Phytothérapie. Paris, France: Maloine; 1992. French.
19. Wichtl M. Herbal Drugs and Phytopharmaceuticals. Boca Raton, FL: CRC Press; 1994.
20. Bartram T. Bartram's Encyclopedia of Herbal Medicine. London, England: Constable and Robinson Ltd; 1998.
21. Weiss R. Herbal Medicine. Beaconsfield, England: Beaconsfield Publishers Ltd; 1988.
22. Chadwick LR, Pauli G, Farnsworth NR. The pharmacology of *Humulus lupulus* L. (hops) with an emphasis on estrogenic properties. Phytomedicine. 2006;13:119 -131.
23. Buckwold VE, Wilson RJ, Nalca A, Beer BB, Voss TG, Turpin JA, et al. Antiviral activity of hop constituents against a series of DNA and RNA viruses. Antiviral Research. 2004;61:57-62.
24. Bauer R. Quality criteria and standardization of phytopharmaceuticals: Can acceptable drugs standards be achieved? Drug Information Journal. 1998;32:101-110.
25. Meier B. The extraction strength of ethanol/water mixtures commonly used for the processing of herbal drugs. Planta Med. 1991;57(S2):A26.
26. Wendakoon C, Calderon P, Gagnon D. Evaluation of Selected Medicinal Plants Extracted in Different Ethanol Concentrations for Antibacterial Activity against Human Pathogens. Journal of Medicinally Active Plants. 2012;1(2):60-68. Accessed 2 February 2013. Available at: <http://scholarworks.umass.edu/jmap/vol1/iss2/4>.
27. Munsel VK, Huber K. Investigation on the suitability various extraction verfahren zur Herstellung eines Kammillenfluid extraktes. Pharmaceutica Acta Helvetiae. 1961;36:194-204. German.
28. Balansard G, Chausse D, Boukef K, Toumi A, Jadot G. Criteria for choosing an extract of Viburnum, *Viburnum prunifolium* L., in function of its toniveineuse and spasmolytic actions. Medicinal plants and herbal medicine. 1983;17(3):123-132. French

© 2014 Gagnon et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history.php?iid=608&id=13&aid=5411>