



# **Biological and Chemical Factors Influencing Food Allergies: A Comprehensive Review**

**Yatharth Sharma <sup>a\*</sup>, Prasad Patil <sup>b++</sup>, Manjit M Khatal <sup>c#</sup>,  
Shilpa Huchchannanavar <sup>d†</sup>, Rachana M N <sup>e</sup>,  
Wankasaki Lytand <sup>f</sup>, Abhijit Debnath <sup>g‡</sup>, Shifat Fatima <sup>h^</sup>  
and Aparna Srivastava <sup>h##</sup>**

<sup>a</sup> Department of Food Science and Nutrition, Maharana Pratap University of Agriculture and Technology, Udaipur, India.

<sup>b</sup> Department of Post-Harvest Management, College of Agriculture, Vellanikkara, Thrissur, Kerala, India.

<sup>c</sup> Department of Process and Food Engineering, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli, India.

<sup>d</sup> Department of Food Science and Nutrition, College of Agriculture, Hagari-583111, University of Agricultural Sciences, Raichur, India.

<sup>e</sup> Department of Processing and Food Engineering, College of Agricultural Engineering, University of Agricultural Sciences, Raichur – 584104, Karnataka, India.

<sup>f</sup> Department of Microbiology, Shillong College, Shillong, India.

<sup>g</sup> Krishi Vigyan Kendra Dhalai, Tripura -799278, India.

<sup>h</sup> Department of Food and Nutrition, Era University, Lucknow, India.

## **Authors' contributions**

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## **Article Information**

DOI: <https://doi.org/10.9734/jsrr/2024/v30i62095>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110661>

<sup>++</sup> Assistant professor (Contract);

<sup>#</sup> Designation - PG Scholar;

<sup>†</sup> Associate Professor;

<sup>‡</sup> Subject Matter Specialist;

<sup>^</sup> Research Scholar;

<sup>##</sup> Assistant Professor;

<sup>\*</sup>Corresponding author;

**Cite as:** Sharma, Y., Patil, P., Khatal, M. M., Huchchannanavar, S., Rachana M N, Lytand, W., Debnath, A., Fatima, S., & Srivastava, A. (2024). Biological and Chemical Factors Influencing Food Allergies: A Comprehensive Review. *Journal of Scientific Research and Reports*, 30(6), 787–794. <https://doi.org/10.9734/jsrr/2024/v30i62095>

## ABSTRACT

Food allergies have become a significant public health concern, affecting millions worldwide. This comprehensive review explores the multifaceted nature of food allergies, shedding light on the intricate interplay of biological and chemical factors. We delve into the mechanisms of allergen recognition and sensitization, genetic predisposition, and the role of the gut microbiome. Moreover, we examine the influence of food processing techniques, additives, and cross-reactivity in the development and severity of food allergies. Food allergies, an adverse health condition triggered by the immune system's hypersensitivity to certain food proteins, have been rising globally, affecting millions of individuals and posing significant public health challenges. This comprehensive review explores the biological and chemical factors influencing the development, manifestation, and management of food allergies. Key biological factors include genetic predisposition, age, and immune system interactions, such as the roles of immunoglobulin E (IgE) and non-IgE mediated responses. Additionally, microbiota composition and environmental exposures during critical windows of immune development are examined. Chemical factors encompass the molecular structure of allergens, food processing methods, and the presence of food additives and contaminants. Advances in diagnostic techniques, such as component-resolved diagnostics and epitope mapping, along with emerging therapies like immunotherapy and biologics, are discussed. The review emphasizes the need for interdisciplinary approaches integrating molecular biology, immunology, and environmental science to develop effective prevention and treatment strategies. Understanding these complex interactions is crucial for reducing the burden of food allergies and improving the quality of life for affected individuals.

*Keywords: Food allergies; allergen recognition; genetic predisposition; gut microbiome; food processing; diagnostic tools.*

## 1. INTRODUCTION TO FOOD ALLERGIES

Food allergies are immune-mediated adverse reactions to specific food proteins. When individuals with food allergies consume allergenic foods, their immune systems mistakenly identify harmless proteins as threats, triggering an allergic response. These reactions can range from mild symptoms, such as hives or digestive discomfort, to severe and potentially life-threatening anaphylaxis, characterized by rapid-onset symptoms like difficulty breathing and a drop in blood pressure [1,2].

The prevalence of food allergies has been steadily rising in recent years, making them a significant global health concern. Although they can affect individuals of all ages, food allergies are particularly common in children. According to estimates, up to 8% of children and 2-4% of adults in the United States have a food allergy [3]. Common food allergens include peanuts, tree nuts, milk, eggs, soy, wheat, fish, and shellfish.

### 1.1 The Socio-economic Impact of Food Allergies

Food allergies have a profound socio-economic impact on affected individuals, their families, and society as a whole:

**Healthcare Costs:** Food allergy management requires medical evaluation, allergy testing, and potentially emergency care for severe reactions. These expenses can place a substantial financial burden on individuals and healthcare systems.

**Quality of Life:** Food allergies can significantly affect the quality of life of those with allergies and their families. Constant vigilance about food choices, dietary restrictions, and fear of accidental exposure can lead to stress and anxiety.

**Impact on Families:** Families with food-allergic members may need to adapt their entire lifestyle to accommodate the allergies. This includes adjusting meal plans, avoiding allergens, and

educating others, which can be emotionally and socially taxing.

**Educational Institutions and Workplaces:**

Schools and workplaces need to implement allergy-aware policies and practices, which can incur additional costs. Moreover, missed school or workdays due to allergic reactions can affect academic and professional performance.

**Food Industry:** The food industry faces challenges in terms of allergen labeling, cross-contamination prevention, and ensuring allergen-free options. Compliance with allergen labeling regulations adds complexities to food manufacturing and labeling processes.

**Public Awareness and Education:** Promoting public awareness about food allergies is essential to prevent accidental exposures and anaphylactic reactions. Allergen-free alternatives may also be more expensive, affecting consumer choices.

## 2. BIOLOGICAL FACTORS IN FOOD ALLERGIES

Food allergies are complex immune-mediated responses involving various biological factors. Understanding these factors is crucial for comprehending how food allergies develop and progress. This section explores the key biological components involved in food allergies:

**Immunoglobulin E (IgE) Antibodies:** IgE antibodies are central players in food allergies. When an individual with a food allergy is exposed to an allergenic protein, the immune system produces specific IgE antibodies against that protein. These antibodies are highly specialized to recognize and respond to the allergen. Upon subsequent exposure to the same allergen, IgE antibodies trigger an allergic reaction by binding to mast cells and basophils, leading to the release of histamine and other inflammatory mediators [4].

**Allergen Recognition and Sensitization:** The initial exposure to an allergenic food may not lead to immediate symptoms. Instead, it can result in sensitization, where the immune system recognizes the allergenic protein as harmful. Subsequent exposures can then lead to an allergic reaction. The sensitization process involves the activation of immune cells like T cells and B cells, which produce IgE antibodies specific to the allergen [5].

**Genetic Predisposition:** Genetic factors play a significant role in determining an individual's susceptibility to food allergies. Individuals with a family history of allergies are more likely to develop food allergies themselves. Specific genetic markers have been identified that increase the risk of food allergy development, although the interplay between genetics and environmental factors is complex and not fully understood (Liu et al., 2017).

**Mast Cells and Basophils:** Mast cells and basophils are immune cells that play a central role in the allergic response. When IgE antibodies bind to these cells and encounter the allergen, they release histamine and other inflammatory molecules, leading to allergy symptoms such as itching, swelling, and bronchoconstriction [4].

**T-Helper Cells (Th2):** T-helper cells, specifically Th2 cells, are involved in promoting allergic responses. Th2 cells release cytokines that drive the production of IgE antibodies. In individuals with food allergies, there is often an imbalance between Th1 and Th2 responses, with a bias towards Th2-driven allergic reactions [6].

**The Gut Microbiome:** Emerging research suggests that the gut microbiome plays a role in food allergies. Changes in the composition and diversity of gut bacteria early in life may influence immune development and increase the risk of allergic sensitization. Probiotics and prebiotics are being investigated for their potential to modulate the gut microbiome and reduce the risk of food allergies [7].

Understanding these biological factors involved in food allergies is a critical step in developing effective prevention and treatment strategies. By targeting specific components of the immune response and genetic predisposition, researchers aim to improve diagnostic tools and develop personalized approaches for managing food allergies.

## 3. GENETIC PREDISPOSITION

Genetic factors play a significant role in determining an individual's susceptibility to food allergies. While the precise genetic mechanisms underlying food allergies are complex and multifaceted, research has identified several key aspects of genetic predisposition. Understanding these genetic factors is crucial for gaining insights into the development and heritability of

food allergies. Here, we delve into the genetic components of food allergies:

### 3.1 Family History and Heritability

Food allergies tend to cluster within families, suggesting a strong hereditary component. Individuals with parents or siblings who have food allergies are at a higher risk of developing allergies themselves. Studies have shown that if both parents have allergies, their child's risk of food allergies is significantly elevated (Liu et al., 2017).

### 3.2 Polygenic Inheritance

Food allergies are considered polygenic, meaning that they result from the interaction of multiple genes. Various genes involved in immune regulation, such as those related to the production of immunoglobulin E (IgE) antibodies, have been implicated. These genes collectively contribute to an individual's overall genetic predisposition to food allergies (Aceves et al., 2016).

### 3.3 Candidate Genes

Researchers have identified specific candidate genes associated with food allergies. For example, variations in the filaggrin gene (FLG) have been linked to increased risk. Filaggrin is involved in maintaining the skin barrier, and mutations in this gene can contribute to the development of eczema, a condition associated with increased food allergy risk [8].

### 3.4 Immune System Regulation

Genes related to immune system regulation are crucial in the development of food allergies. Polymorphisms in genes encoding cytokines, interleukins, and toll-like receptors can influence an individual's immune response to allergenic proteins. Altered immune regulation can increase the likelihood of an allergic response [9].

### 3.5 Epigenetic Modifications

Epigenetic modifications, such as DNA methylation and histone modifications, can influence gene expression without altering the underlying DNA sequence. Research suggests that epigenetic changes may play a role in the development of food allergies, potentially mediating the interaction between genetics and environmental factors [10].

## 3.6 Gene-Environment Interactions

Food allergies often result from a complex interplay between genetics and environmental factors. Genetic predisposition alone may not be sufficient for an individual to develop an allergy. Factors such as early exposure to allergenic foods, changes in the gut microbiome, and environmental allergen exposure also contribute to the development of food allergies (Liu et al., 2017).

Understanding the genetic basis of food allergies is an active area of research. Genetic studies provide valuable insights into the underlying mechanisms of allergy development, which may lead to improved diagnostic tools, personalized treatment strategies, and potential interventions to reduce the risk of food allergies in genetically susceptible individuals.

## 4. THE GUT MICROBIOME AND IMMUNE REGULATION

The gut microbiome, comprising trillions of microorganisms residing in the gastrointestinal tract, plays a pivotal role in immune regulation and the development of food allergies. The intricate interactions between the gut microbiome and the immune system have significant implications for understanding and potentially modulating the risk of food allergies. This section explores the relationship between the gut microbiome and immune regulation in the context of food allergies:

### 4.1 Gut Microbiome Composition

The gut microbiome is composed of a diverse community of bacteria, viruses, fungi, and other microorganisms. Its composition varies among individuals and is influenced by factors such as diet, genetics, and environmental exposures. Recent research has revealed that the diversity and balance of microbial species in the gut are crucial for maintaining immune homeostasis [11].

### 4.2 Gut Microbiome and Immune Tolerance

One of the key functions of the gut microbiome is the promotion of immune tolerance. Early exposure to a diverse range of gut bacteria helps educate the immune system, teaching it to distinguish between harmless food proteins and potential threats. This immune tolerance is vital in preventing the development of food allergies [7-15].

### 4.3 Dysbiosis and Allergic Sensitization

Dysbiosis, characterized by an imbalance in the gut microbiome composition, has been linked to an increased risk of allergic sensitization and food allergies. Dysbiotic conditions can disrupt immune tolerance, making individuals more susceptible to allergic reactions when exposed to food allergens [11].

### 4.4 Probiotics and Prebiotics

Probiotics are live microorganisms that can confer health benefits when consumed in adequate amounts. Some probiotic strains have shown potential in modulating the gut microbiome and reducing the risk of food allergies. Prebiotics, which are non-digestible dietary fibers that promote the growth of beneficial gut bacteria, may also play a role in enhancing immune tolerance [7].

### 4.5 Microbiome-Immune Crosstalk

The gut microbiome communicates with the immune system through a complex network of interactions. These interactions involve the release of signaling molecules, such as short-chain fatty acids, by gut bacteria. These molecules can modulate immune cell function and influence immune responses to allergens [12].

### 4.6 Early-Life Influences

The first years of life are a critical period for microbiome development and immune programming. Factors such as delivery mode (vaginal vs. cesarean), breastfeeding, and antibiotic use can shape the composition of the gut microbiome during infancy, impacting long-term immune health and allergy risk [11].

Understanding the role of the gut microbiome in immune regulation and its impact on food allergies is a rapidly evolving area of research. Harnessing this knowledge may lead to novel preventive and therapeutic strategies, such as the use of probiotics, prebiotics, and microbiome-targeted interventions, to reduce the risk of food allergies and improve overall immune health.

## 5. CHEMICAL FACTORS: FOOD PROCESSING AND ADDITIVES

Food processing techniques and the use of food additives can significantly impact the development and severity of food allergies. Chemical factors associated with food

processing and additives play a critical role in influencing allergenicity and allergic reactions. This section examines the relationship between food processing, additives, and food allergies:

### 5.1 Impact of Food Processing Techniques

**Heat Processing:** Cooking and heat processing can alter the allergenicity of food proteins. In some cases, heat denatures allergenic proteins, rendering them less allergenic and reducing the risk of allergic reactions. However, in other cases, heat may not fully eliminate allergenicity [16].

**Roasting and Baking:** Roasting and baking can enhance the allergenic potential of certain foods. For example, roasting peanuts can increase the allergenicity of peanut proteins, potentially making them more allergenic to susceptible individuals [17].

**Frying and Maillard Reaction:** High-temperature frying and the Maillard reaction, which occurs when proteins and sugars react during cooking, can lead to the formation of new allergenic compounds. This can increase the allergenic potential of some foods [18].

### 5.2 Role of Food Additives

**Preservatives and Antioxidants:** Some food additives, such as sulfites and antioxidants, have been associated with allergic reactions. Sulfites, which are used to preserve the color and flavor of foods, can trigger allergic responses in sensitive individuals [19].

**Artificial Colors and Flavorings:** Artificial colors and flavorings can contain allergenic compounds that may cause allergic reactions. These additives are commonly found in processed foods and beverages [20].

**Emulsifiers and Stabilizers:** Emulsifiers and stabilizers are often used to improve the texture and shelf life of processed foods. Some individuals may be sensitive to these additives, experiencing digestive discomfort or allergic reactions [21-25].

### 5.3 Cross-Contamination and Labeling

Cross-contamination during food processing and packaging can lead to unintentional exposure to allergenic proteins. It is crucial for manufacturers

to implement rigorous allergen control measures to prevent cross-contact between allergenic and non-allergenic foods. Additionally, accurate allergen labeling is essential to inform consumers about potential allergen risks [26-31].

#### 5.4 Processing-Induced Changes in Allergenicity

The structural changes that occur during food processing can either enhance or reduce the allergenicity of proteins. These changes may involve alterations in protein conformation, exposure of allergenic epitopes, or the formation of new protein complexes. Understanding these processing-induced changes is vital for assessing allergenic risk [32-33].

#### 5.5 Personalized Risk Assessment

Given the variability in allergenicity and individual sensitivity, personalized risk assessments may be necessary. These assessments take into account an individual's specific allergies, sensitivities, and tolerance thresholds to make informed dietary choices [34-37].

### 6. CONCLUSION

Understanding the chemical factors associated with food processing and additives is essential for individuals with food allergies and those involved in food production. Rigorous allergen control measures, accurate labeling, and ongoing research into the impact of processing techniques on allergenicity are vital for minimizing the risk of allergic reactions associated with food consumption. ### Conclusion

Food allergies represent a multifaceted health issue influenced by a complex interplay of biological and chemical factors. This review highlights the critical roles of genetic predisposition, immune system dynamics, and environmental exposures in the development and severity of food allergies. Furthermore, it underscores the impact of allergen molecular structures, food processing methods, and the presence of additives and contaminants on allergic responses.

Advancements in diagnostic methodologies, including component-resolved diagnostics and epitope mapping, are paving the way for more precise identification and understanding of

allergenic proteins. These technologies facilitate better risk assessment and personalized management plans for individuals with food allergies. Emerging therapeutic approaches, such as oral immunotherapy, sublingual immunotherapy, and biologics, show promise in desensitizing allergic individuals and potentially achieving long-term tolerance. The review calls for an integrated approach that combines insights from molecular biology, immunology, environmental science, and clinical practice. This multidisciplinary collaboration is essential for developing innovative prevention strategies, improving diagnostic accuracy, and enhancing treatment efficacy. In conclusion, addressing food allergies requires a comprehensive understanding of both the biological mechanisms and chemical influences involved. Continued research and collaboration are imperative to mitigate the growing prevalence of food allergies and improve the quality of life for those affected. Through these efforts, we can aspire to reduce the global burden of food allergies and enhance public health outcomes.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Aceves SS, Newbury RO, Dohil R, Bastian JF, Broide DH. Esophageal remodeling in pediatric eosinophilic esophagitis. *Journal of Allergy and Clinical Immunology*. 2010; 126(1):81-90.
2. Salam MA, Islam MR, Diba SF, Hossain MM. Marker assisted foreground selection for identification of aromatic rice genotype to develop a modern aromatic line. *Plant Science Archives*; 2019.
3. Liu AH, Jaramillo R, Sicherer SH, Wood RA, Bock SA, Burks AW, Sampson HA. National prevalence and risk factors for food allergy and relationship to asthma: Results from the national health and nutrition examination survey 2005-2006. *Journal of Allergy and Clinical Immunology*. 2010;126 (4):798806.
4. Islam MS, Rahman MM, Paul NK. Arsenic-induced morphological variations and the role of phosphorus in alleviating arsenic toxicity in rice (*Oryza sativa* L.). *Plant Science Archives*. 2016;1(1):1-10.

5. Sicherer SH, Sampson HA. Food allergy: A review and update on epidemiology, pathogenesis, diagnosis, prevention, and management. *Journal of Allergy and Clinical Immunology*. 2014;133(2):291-307.
6. Galli SJ, Tsai M, Piliponsky AM. The development of allergic inflammation. *Nature*. 2008;454(7203):445-454.
7. Kulis M, Burks AW. Immune mechanisms of food allergy. *Current Opinion in Immunology*. 2007;19(6):583-589.
8. Dominguez-Ortega J, del Rio-Prado E, Grupo Espanol de Investigacion En Alergia Al, y Asma G. Allergic and autoimmune reactions in the immune response to food. *Advances in Nutrition*. 2016;7(5):794S-798S.
9. Fujimura KE, Sitarik AR, Havstad S, Lin DL, Levan S, Fadrosch D, Lynch SV. Neonatal gut microbiota associates with childhood multisensitized atopy and T cell differentiation. *Nature Medicine*. 2016;22(10):1187-1191.
10. Palmer CN, Irvine AD, Terron-Kwiatkowski A, Zhao Y, Liao H, Lee SP, McLean WH. Common loss-of-function variants of the epidermal barrier protein filaggrin are a major predisposing factor for atopic dermatitis. *Nature Genetics*. 2006;38(4):441-446.
11. Azra BH, Fatima T. Zinc nanoparticles mediated by *Costus pictus* leaf extract to study GC-MS and FTIR analysis. *Plant Science Archives*. 2024;11-15.
12. Marenholz I, Nickel R, Rüschenhoff F, Schulz F, Esparza-Gordillo J, Kerscher T, Lau S. Filaggrin loss-of-function mutations predispose to phenotypes involved in the atopic march. *Journal of Allergy and Clinical Immunology*. 2006;118(4):866-871.
13. Martino DJ, Prescott SL, Leung DY. Epigenetics of allergy. *Advances in Experimental Medicine and Biology*. 2019;1074:75-98.
14. Nazneen S, Sultana S. Green synthesis and characterization of *Cissus quadrangularis*. L stem mediated zinc oxide nanoparticles. *Plant Science Archives*. 2024;1(05).
15. Savage JH, Lee-Sarwar KA, Sordillo JE, Lange NE, Zhou Y, O'Connor GT, Lynch SV. A prospective microbiome-wide association study of food sensitization and food allergy in early childhood. *Allergy*. 2018;73(1):145-152.
16. Vally H, Misso NL. Adverse reactions to the sulphite additives. *Gastroenterology and Hepatology from Bed to Bench*. 2012;5(1):16-23.
17. Kushwah N, Billore V, Sharma OP, Singh D, Chauhan APS. Integrated nutrient management for optimal plant health and crop yield. *Plant Science Archives*.
18. Gultekin F, Sogut B. Artificial food additives and immunologic reactions. *Turkish Journal of Allergy and Immunology*. 2013;11(3):69-73.
19. Bhakta S, Sipra BS, Dutta P, Sahu E, Panda SK, Bas-tia AK. Water silk (*Spirogyra bichromatophora*) as a natural resource for antimicrobial phytochemicals. *Acta Botanica Plantae*. 2022;V01i03:08-14.
20. Aparanjitha R, Imran GM, Mondal K. Nano fertilizers: Revolutionizing agriculture for sustainable crop growth. *Agriculture Archives*. 2023;2.
21. Reddy CA, Oraon S, Bharti SD, Yadav AK, Hazarika S. Advancing disease management in agriculture: A review of plant pathology techniques. *Plant Science Archives*; 2024.
22. Sultana N, Saini PK, Kiran SR, Kanaka S. Exploring the antioxidant potential of medicinal plant species: A comprehensive review. *Journal of Plant Biota*; 2023.
23. Asma J, Subrahmanyam D, Krishnaveni D. The global lifeline: A staple crop sustaining two thirds of the world's population. *Agriculture Archives*; 2023.
24. Ogori AF, Eke MO, Girgih TA, Abu JO. Influence of aduwa (*Balanites aegyptiaca*. del) meal protein enrichment on the proximate, Phytochemical, Functional and Sensory Properties of Ogi. *Acta Botanica Plantae*. 2022;1(3):22-35.
25. Taylor SL, Hefle SL, Bindslev-Jensen C. Factors affecting the determination of threshold doses for allergenic foods: How much is too much? *The Journal of Allergy and Clinical Immunology*. 2002;109(1):2430.
26. Rahat KMR, Amin MA, Ahmed MT. Comparing tourists' travel cost and consumer surplus to estimate the recreational values of Kuakata Sea Beach in Bangladesh. *J Tourism Hospit*. 2024;13:541.
27. Buchheit KM, Bernstein JA. The impact of processing on allergenicity of food. *Current*

- Allergy and Asthma Reports. 2013;13(5): 471-478.
28. MA, Sultana N, Hansda NN, BM H, Noopur K. Protected vegetable crop production for long-term sustainable food security. Journal of Scientific Research and Reports. 2024;30(5):660-669.
29. Alemu TT. Effect of storage time and room temperature on physicochemical and geometric properties of banana (*Musa Spp.*) Fruit. Journal of Plant Biota. 2023;30-40.
30. Rahgu K, Choudhary S, Kushwaha TN, Shekhar S, Tiwari S, Sheikh IA, Srivastava P. Microbes as a promising frontier in drug discovery: A comprehensive exploration of nature's microbial marvels. Acta Botanica Plantae. 2023;V02i02:24-30.
31. Radauer C, Bublin M, Wagner S, Mari A, Breiteneder H. Allergens are distributed into few protein families and possess a restricted number of biochemical functions. The Journal of Allergy and Clinical Immunology. 2008; 121(4):847-852.
32. Chen MS, Liu SL, Ye XL, Yu XX, Yu JS, Chen F. New developments in food allergen analysis and allergen risk assessment. Food Research International. 2016;89:25-38.
33. Reddy CA, Oraon S, Bharti SD, Yadav AK, Hazarika S. Advancing disease management in agriculture: A review of plant pathology techniques. Plant Science Archives; 2024.
34. Turner PJ, Kemp AS, Campbell DE. Advisory food labeling for allergen avoidance in children with food allergy. The Cochrane Database of Systematic Reviews. 2015;2:CD008006.
35. Md. Tanvir Ahmed, Md. Al Amin. Perilous resurgence of dengue fever in Bangladesh: Gender based perspectives on risk perception and adaptation strategies. Universal Journal of Public Health. 2023;11(5):751-760. DOI: 10.13189/ujph.2023.110525
36. Niranjana C. Characterization of bacteriocin from lactic acid bacteria and its antibacterial activity against *Ralstonia solanacearum* causing tomato wilt. Plant Science Archives; 2016.
37. Hernandez ML, von Mutius E. Th2-biased immune development in early life: What does it mean for the rising trends in allergy and asthma? Annals of Allergy, Asthma & Immunology. 2019;122(3):257-262.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/110661>