



## **Report of Antibiotic Resistance in Urban and Rural Wastewaters from West Bengal, India**

**Meesha Singh <sup>a</sup>, Rupsha Karmakar <sup>b</sup>, Sayak Ganguli <sup>c\*</sup>  
and Mahashweta Mitra Ghosh <sup>a</sup>**

<sup>a</sup> Department of Microbiology, St. Xavier's College, (Autonomous); Kolkata, India.

<sup>b</sup> Department of Zoology, Vivekananda College for Women, Kolkata, India.

<sup>c</sup> Department of Biotechnology, St. Xavier's College, (Autonomous); Kolkata, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. Author MS performed the wet lab experiments, authors RK and SG was involved in computational analysis, authors SG and MMG conceptualised the study and evaluated the manuscript. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JPRI/2021/v33i53A33660

#### Editor(s):

(1) Dr. Ana Cláudia Coelho, University of Trás-os-Montes and Alto Douro, Portugal.

#### Reviewers:

(1) E. Kavitha, Sri Venkateshwara Medical College Hospital and Research Centre, India.

(2) Ahmed Sadeq Abou El-Yazed Aresha, Menoufia University, Egypt.

(3) Khaldi Amina, Université sidi bel abbes Algeria, Algeria.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:  
<https://www.sdiarticle5.com/review-history/77759>

**Original Research Article**

**Received 20 September 2021**

**Accepted 30 November 2021**

**Published 04 December 2021**

### **ABSTRACT**

**Aims:** This study aims at comparative identification of antibiotic resistance patterns in bacteria isolated from samples collected from rural environment (LS) and urban environments (SS). Metagenomic profiling gave us insights into the microbial abundance of the two samples. This study focused on culture-based methods for complete identification of antibiotic resistant isolates and estimation of comparative antibiotic resistance among the two samples.

**Study Design:** Untreated medical waste and anthropogenic waste disposal can lead to the propagation of different antibiotic resistant strains in wastewater environments both in urban and rural set ups which provide an insight towards this study approach mentioned in the methodology segment.

**Place and Duration of Study:** Sewer system of a medical facility located in Purulia, India was the collection site for liquid sludge. Solid sludge and associated wastewater were collected in vicinity of a large urban medical facility from central Kolkata, India.

**Methodology:** Physico-chemical properties were analyzed followed by microbiological and biochemical characterization. The antibiotic resistance patterns were determined by Kirby-Bauer disc diffusion assay. Potent multidrug resistant isolates were identified using 16srRNA gene amplification followed by Phylogenetic profiling, using CLC Genomics workbench.

**Results:** We observed maximum resistance in an *E. coli* isolate which was resistant up to 22 antibiotics. Combined data for resistance from urban and rural samples were found to exhibit 83.9% resistance to beta lactams, 85.7% to macrolides, 44.2% to fluoroquinolones, 50% to glycopeptides and cephalosporins, 35.7 % to carbapenems and sulfonamides, 28.5 % to tetracycline, and 23.8 % to aminoglycosides.

**Conclusion:** The high prevalence of antibiotic-resistant bacteria harbouring diverse resistance traits across samples indicated towards probable horizontal gene transfer across environmental niches. This study can prove to be useful to understand and map the patterns of resistance and stringently apply the counter measures related to public health practices.

**Keywords:** Antibiotic resistance; ARB; environmental resistome; antibiotic pollution; wastewater; Sludge.

## 1. INTRODUCTION

The wonder molecules which were once used as lifesaving drugs have gradually been rendered ineffective by the evolution of microorganisms which have devised mechanisms of resistance against them. Extensive use of antibiotics in healthcare and veterinary sectors along with their continuous unmonitored discharge in natural environments have successfully built up a reservoir of antibiotic resistant bacteria (ARB) which co evolve by acquisition of antibiotic resistance genes (ARG). This has led the world health organization to identify pathogenic strains of microorganisms which are potential superbugs exhibiting patterns of multidrug resistance thus directly diminishing the therapeutic potential of the antibiotics [1].

In India there is tendency of overuse of broad-spectrum penicillins owing to their availability and low cost. As a result, recent reports suggest that it is the leading country in per person antibiotic consumption which automatically results in the buildup of drug resistant pathogens [2] leading to the coinage of the phrase 'Antimicrobial resistance (AMR) capital of the world'[3].

There are several underlying factors such as inappropriate disposal of antibiotics, misuse etc. which cause intake and transfer of antibiotic resistant gene/genes in resistant bacteria [4]. The load of antibiotic resistance genes in natural environments contributes towards efficient transfer of these gene modules or collection of antibiotic resistant genes to different bacterial strains using mechanisms such as horizontal gene transfer, recombination or mutations under the selection pressure of different pollutants such

as pesticides and heavy metals [5]. Some bacteria have been reported to utilize the antibiotic residues present as potential carbon source [6]. In ecological niches, the resistant genes are pooled together and as a result of this, there is an uptake of resistant genes among the bacterial neighbours horizontally. The lack of record of contamination is evident from the current reports of bacterial resistance towards third generation beta lactams and fourth generation cephalosporins [7]. There have been few reports on a lesser-known phenomenon, "environmental resistance loop" which refers to the transfer of resistant bacteria and antibiotic residues from wastewater treatment plants to the riverine systems and agricultural lands, finally reaching back to human beings and animals [8]. There is a growing concern that this phenomenon could render current and future antibiotics ineffective. According to current literatures, the mobile resistant genetic elements such as insertion sequences, transposons, integrons, and plasmids from clinical samples have been detected in hospital associated wastewaters, within a very minimal time interval following their report in hospitals [9,10]. The unique characteristics of resistome are quite evident and thus there is a dire need for an integrated approach which includes effective wastewater treatment and continuous monitoring of the resistome. The main anthropogenic sources of dissemination of ARBs are effluents from wastewater and hospital discharge where horizontal gene transfer is very dominant [11] and as a result, the "difficult to treat" infections are increasing globally [12]. The causative agents and their representative antibiotic resistance profiles are present in low percentages in wastewaters, as compared to that

in clinical settings. However, this presents a serious public health issue.

We still are in the nascent stage of our understanding regarding the types and prevalence of antibiotic resistance in the environment. Though, the standardization and organization of antibiotic resistance data of clinical origin has been curated, information regarding resistance of environmental bacteria is still very fragmented [13]. In India, according to the Resistance map resource [14] microbes resistant to aminoglycosides, carbapenems and cephalosporins pose the highest threat and have shown a steady pattern of increase from 2008 to 2020. The current pandemic has also opened a Pandora's box in terms of antibiotic usage, recent data from five different countries, suggests that there are associated bacterial infections in 6-9% of COVID-19 diagnoses of which 3-5% have been diagnosed during initial hospitalization while 14-3% post recovery. This trend is higher for patients who have received intensive critical care thus nosocomial infections cannot be ruled out [15]. This multicenter US based study also reported that almost 72% of the COVID patients were treated with antibiotics, even when there was no absolute clinical necessity [15]. The status of use of antibiotics in less developed countries are so scratchy that it is difficult to understand the actual burden that is being added to the environment as each day passes. Researchers believe that antimicrobial resistance might become even worse after COVID-19 due to their excessive use of antibiotics in humans, misuse in agriculture, and the unavailability of new formulations in the pipeline.

The aim of this study was to identify the diverse antibiotic resistant bacteria profiles in environmental wastewaters with respect to multiple antibiotics from diverse classifications. In a previous study the bacterial abundances of these two sites were reported using comparative metagenomics [16]. The metagenomic profiling gave us insights into the microbial abundance of the two samples. This study focused on culture-based methods for complete identification of antibiotic resistant isolates and estimation of comparative antibiotic resistance among the two samples.

## 2. MATERIALS AND METHODS

Wastewater environments were initially identified for their proximity to large medical facilities and then sample was collected as described in the

previous report [16]. The microbiological culturing was initiated within 10 hours of collection. Stock samples were stored at 4°C for further processing.

### 2.1 Study Area

The first area of this study is Purulia. Its latitudinal and longitudinal extents are from 22°42'35'' to 23°42'00'' North and from 85°49'25'' to 86°54'37'' East. The region is climatically characterised as a dry and arid zone with parts of the Chotanagpur plateau region contributing towards its warm and humid conditions. The sample collection site was wastewater effluent collected in vicinity of a medical facility located in Purulia. The climatic conditions were recorded; Temperature: 42°C, Relative humidity: 68%, Time of collection: 12 pm.

The second area of this study, Kolkata, is located in the eastern part of India. The Calcutta Municipal Corporation has an area of 185 sq km in total. A large part of the city historically was marshy wetlands, remnants of which can still be found especially towards the eastern parts of the city [17]. The sample collection site was solid sludge and associated wastewater collected in vicinity of a large urban medical facility located in central Kolkata. The climatic conditions were recorded; Temperature: 34°C, Relative humidity: 95%, Time of collection: 11:45 am.

### 2.2 Evaluation of Physicochemical Parameters of the Sampling Stations

The physico-chemical parameters of the effluents were analyzed as per standardized EPA protocols [18]. The parameters analyzed were Chemical oxygen demand (COD), Total organic carbon (TOC), Nitrate, Ammonia, total Kjeldahl nitrogen, total Phosphorus and heavy metals, viz. Chromium (Cr), Mercury (Hg), Lead (Pb), Cadmium (Cd), Arsenic (As).

### 2.3 Microbiological Characterization

The effluent samples were serially diluted and plated onto nutrient agar plates (pH 6.6) and were incubated overnight at 37°C. Isolated colonies in higher dilution were observed on each type of plates. These colonies were then sub-cultured in nutrient agar plates for pure colony isolation and subsequently Gram stained. The colonies were observed under 100X Magnification using Bright Field Light Microscope

for morphological identification. The Gram positive and Gram-negative bacterial isolates were then presumptively identified through a series of cultural and biochemical tests according to the criteria described in Bergey's Manual of Determinative Bacteriology [19].

## 2.4 Biochemical Characterization

The isolates were subjected to routine biochemical tests: Catalase, Oxidase, Nitrate reduction, Indole, Methyl red, Voges-Proskauer, Citrate utilization, Urease, Starch hydrolysis, Carbohydrate fermentation [20].

## 2.5 Antibiotic Resistance Profiling

To determine the susceptibility of the isolates towards different antibiotics Kirby-Bauer disc diffusion assay was done. The antibiotic against which the susceptibility of the isolates was tested was selected on the basis of recommendations given in CLSI 2019, antibiotics belonging to diverse classifications and generations were used to check the sensitivity of the isolates. (Supplementary Table B). Disc diffusion assays were done using antibiotic discs from HiMedia Laboratories (Mumbai, India). Sensitivity of the isolates against antibiotics (intermediate or susceptible) was deduced on the basis of the information given in the manufacturer's manual. Results were interpreted on the basis of CLSI guidelines [21]. We classified an isolate to be multi drug resistant (MDR) if it exhibited resistance to antibiotics belonging to three or more classes.

## 2.6 Molecular Identification

Genomic DNA was isolated from different bacterial isolates using standard protocols and 16SrDNA fragment were amplified using universal 16S primers 27f (5'AGAGTTTGATCCTGGCTCAG3') and 1492r (5'TACGGTTACCTGTTCGACTT3') [20], and sequenced. Raw sequences were assembled using the Cap3 Contig Assembly –in silico program. All the sequences were used to identify the bacteria with the help of the BLASTn program and were submitted to GenBank [22].

## 2.7 Phylogenetic Analysis

For constructing the phylogenetic tree, initially two programs were used to generate the multiple sequence alignment for a rapid validation of sequence scores -they were Clustal omega [23]

and Muscle [24]. Based on the consensus multiple sequence alignment, phylogenetic tree was determined with 100 bootstrap replicates [25] using the phylogenetic tree module of the CLC Genomics workbench. Here both neighbour joining-based methods [26] and maximum likelihood [27] based methods were tested to evaluate the tree generated and the best consensus tree was chosen for analysis and interpretation. The tree file was visualized in Fig Tree to distinguish the different clade.

## 3. RESULTS AND DISCUSSION

### 3.1 Physico-Chemical Characterization

The physico-chemical parameters of the effluents were identified with the purpose of observing driving factors behind widespread antibiotic resistance and horizontal gene transfer. Heavy metals such as Cd, Pb, Hg were found to be higher than the permissible limits in both effluents (Fig. 1a). Heavy metals have been reported to be co-selecting agents, which promote resistance by co-resistance and cross-resistance mechanisms [28]. Total Phosphorus was higher than the permissible limits. COD was also found to be higher than the permissible limits (Fig. 1b).

### 3.2 Microbiological Characterization

The total number of isolated bacteria for LS and SS datasets was 19 and 23 respectively. The study was focused on multidrug resistant strains due to it was narrowed down to 14 isolates with 10 isolates from LS dataset and 4 isolates from SS dataset. Among the selected isolated bacteria, 8 were Gram negative rods (LS 2, LS 4, LS 6, LS 9, LS 17, LS 21, SS 7 and SS 19) and 6 were Gram-positive rod-shaped bacteria (LS 12, LS 14, LS 21, LS 23, LS 25 and SS 20).

### 3.3 Biochemical Characterization


The isolated bacteria were subjected to various biochemical tests and the results were tabulated (Supplementary Table A and C).

### 3.4 Antibiotic Resistance Patterns

The antibiotic resistance was determined in terms of zone of inhibition (cm) data and it was found that isolates obtained from Purulia (rural area) was found to be much higher than that in Kolkata (urban area). *E. coli* isolate (LS 2) was resistant against 22 antibiotics out of 24

antibiotics tested (Supplementary Table B) which is in conformation with the public data released by ICMR. The percentage of resistance was higher in *E. coli* isolates compared to other isolates. Some *E. coli* isolates (LS 2, LS 9, LS 17) showed variable pattern of susceptibility which may be attributed to its higher relative abundance in fecal contaminated environments or increased competence towards receiving

resistant gene containing foreign plasmids [29]. A study from diarrhoea affected children in Mexico revealed 73% isolates of *E. coli* 73% resistant to ampicillin [30]. In Pakistan, [31] identified a multi antibiotic resistant isolate of *E. coli* which was resistant to cefotaxime, ceftazidime, gentamycin, ciprofloxacin, imipenem. This study also exhibits similar patterns of resistance in *E. coli*. (Table 1) (Supplementary Table D).



**Fig. 1. Heavy metal concentrations in samples (Graphical representation of the estimation of heavy metal estimation in LS and SS sample sets with parameters on y-axis and concentrations (mg/l) on x-axis) (a) and Physico-chemical parameters of samples (Graphical representation of concentration of various analytes for LS and SS sample sets with parameters on y-axis and concentrations (mg/l) on x-axis) (b)**

**Table 1. Identified isolates with varying antibiotic resistance patterns**

Sample ID	Isolate	Antibiotics resistant pattern
LS 2	<i>Escherichia coli</i>	AMP, CIP, GEN, NOR, CTR, NAL, AK, AMC, Cd, OFX, VAN, ERY, MET, Cf, TET, DO, CTX, IPM, CXM, LE, COT, CX
LS 4	<i>Shigella flexneri</i>	AMP, CIP, NAL, AMC, Cd, E, MET, Cf, CXM
LS 6	<i>Klebsiella pneumoniae</i>	AMP, CIP, GEN, CTR, AK, AMC, OFX, NAL, AMC, Cd, E, MET, Cf, TET, DO, CTX, IPM, CXM, COT
LS 9	<i>Escherichia coli</i>	AMP, NOR, NAL, Cd, MET, Cf
LS 12	<i>Bacillus safensis</i>	AMP, Cd, ERY, MET, Cf, CXM
LS 14	<i>Bacillus australimaris</i>	AMP, NAL, AMC, Cd, VAN, ERY, MET, Cf, CX
LS 17	<i>Escherichia coli</i>	AMP, NAL, Cd, OFX, VAN, ERY, MET
LS 21	<i>Comamonas aquatica</i>	AMP, GEN, NOR, CTR, NAL, AK, AMC, Cd, OFX, ERY, MET, Cf, TET, DO, CTX, IPM, CXM
LS 23	<i>Lysinibacillus fusiformis</i>	AMP, CIP, GEN, NOR, CTR, NAL, AK, AMC, Cd, OFX, ERY, MET, Cf, TET, CTX, IPM, CXM, COT
LS 25	<i>Oceanobacillus caeni</i>	AMP, GEN, NOR, CTR, NAL, AK, AMC, Cd, OFX, VAN, ERY, MET, Cf, CTX, IPM, CXM, COT
SS 3	<i>Terribacillus halophilus</i>	AMP, VAN, ERY, Cf
SS 7	<i>Citrobacter freundii</i>	AMP, CIP, CTR, NAL, Cd, OFX, ERY, MET, Cf, TET
SS 19	<i>Comamonas aquatica</i>	AMP, NAL, Cd, VAN, MET
SS 20	<i>Bacillus pumilus</i>	AMP, CIP, CTR, NAL, AMC, Cd, MET, Cf, CXM

*K. pneumoniae* is one of the commonest isolates in both hospital and community acquired infections [32]. MDR and carbapenem resistant *K. pneumoniae* has become a major therapeutic challenging scenario in several countries due to the lack of alternative existing antibiotics. Manikandan and Amsath [33] observed high degree of resistance in *K. pneumoniae* isolated from a urine sample. They reported resistance to ampicillin, ceftazidime, cefotaxime, ciprofloxacin and gentamicin. In our study, *K. pneumoniae* isolate (LS 4) also exhibits resistance against beta lactams, fluoroquinolones, aminoglycoside, macrolides, tetracycline, cephalosporins, carbapenems and sulphonamide.

Several reports from around the world specially, India, Canada, Israel, Argentina, Turkey, Lebanon, Iran, China, Japan and South Korea, have reported *Shigella* spp. harbouring different types of ESBL genes [34]. In our study we found that *Shigella flexneri* isolate (LS 2) showed variable resistance towards beta lactams, fluoroquinolones, macrolides and cephalosporins.

Several strains of the commonly occurring *Lysinibacillus fusiformis* have been isolated from multiple environments including agriculture soil and factory wastewater [35]. However, there are no reports corresponding to the emergence of multidrug resistance in it. *L. fusiformis* has been reported to be sensitive to tetracycline previously [36]. In our study, *L. fusiformis* (LS 23) was found to be resistant against beta lactams, fluoroquinolones, macrolides, tetracycline, cephalosporins and sulphonamide.

Bacteria of the *Bacillus pumilus* clade are ubiquitous in a wide variety of terrestrial and marine environments, ranging from stratospheric air to deep-sea sediments and from soil to living beings [37,38, 39]. The following species belonging to *B. pumilus* clade: *B. pumilus* (LS 16), *B. safensis* (LS 21) and *B. australimaris* (LS 14) were identified in this study. The antibiotic resistance patterns of the sister clade isolates are quite similar in nature with resistance shown against beta lactams, fluoroquinolones, macrolide and cephalosporins. This presents an interesting insight into the evolutionary link of antibiotic resistance genes across various species.

*Comamonas* species are occasional human pathogens found in contaminated environments [40]. Their identification has been challenging

with several laboratories ending up reporting them as non fermentative gram-negative bacilli that could not be further identified [41]. We have identified two *C. aquatica*, one each from both the samples (LS 17 and SS 20), which show variable resistance against beta lactams, fluoroquinolones and glycopeptide class of antibiotics. This is probably the first report of antibiotic resistance in *Comamonas* species from a waste water environment.

*Oceanobacillus caeni* was first isolated in South Korea as a component of activated sludge in a Bacillus-dominated wastewater treatment plant [42]. In our study the isolate (LS 22) was obtained from rural wastewater and it showed variable resistance against beta lactams, fluoroquinolones, macrolides, cephalosporins and sulphonamide. The pathology and antibiotic resistance patterns of this species have not been conclusively studied before.

*Citrobacter freundii* is a frequent cause of nosocomial infections and a known cause of diarrheal infections, has increasingly become multidrug resistant [43]. *C. freundii* isolate (LS 25) was identified and showed resistance against beta lactams, fluoroquinolones, aminoglycoside, macrolides, cephalosporins and tetracycline.

*Terribacillus halophilus* was originally isolated from field soil in Japan [44]. The isolate (SS 3) showed variable antibiotic resistance patterns against beta lactams, glycopeptides, macrolide and cephalosporin. This is the first report of occurrence of antibiotic resistance in this species. (Fig. 2a and b).

This study identifies 83.9% of the isolates to exhibit resistance to beta lactams with 100% resistance to Ampicillin [45]. Reported higher probability of occurrence of ESBL-producing *K. pneumoniae* and *E. coli* isolates in hospital effluent, Waste Water Treatment Plants (WWTP) and river samples, respectively whereas, hospital effluent, sanitary effluent, outflow sewage and surface water samples were richer in quinolone resistant isolates. In this study, high macrolide resistance among the isolates (85.7%) which clearly points to the diverse antibiotic resistance potential of the effluents. The resistance to fluoroquinolones was 44.2 %, 50% for glycopeptides and cephalosporins, 35.7 % for carbapenems and sulfonamides, 28.5 % for tetracycline, 23.8% for aminoglycosides. All the isolates were however, found to be susceptible to chloramphenicol and streptomycin which may be


due to the fact that these antibiotics are prescribed very sparsely by healthcare professionals and medical practitioners in the vicinity of the collection spots.

Phylogenetic profiling revealed the presence of 4 distinct phylogenetic clusters (Table 2) (Fig. 3). The largest cluster had 5 individual OTUs (LS 4, SS 7, LS 17, LS 9, SS 19). In this group, all the isolates showed resistance against AMP, NAL, Cd, VAN and MET antibiotics. It is interesting to note that this group contains isolates obtained from different geographical and sampling conditions which indicate the possibility of genetic exchange amongst allochthonous species [46]. The cluster containing isolates LS 6, LS 2, LS 21 show resistance against a wide range of antibiotics such as beta lactams, fluoroquinolones, aminoglycoside, macrolides, tetracycline, cephalosporins, carbapenems and sulphonamides, evolving probably through Darwinian forces [47]. The sister group with OTUs obtained from different sampling conditions such as LS 14, SS 20, LS 12 show resistance against beta lactams and cephalosporins both groups inhibiting bacterial cell wall synthesis. The cluster with OTUs LS 23, SS 3 and LS 25 contains halophilic organisms with resistance against some common antibiotics such as beta lactams and cephalosporins. LS 2 and LS 6 belong to the sister clades and exhibit similar antibiotic resistance patterns viz. resistance against beta lactams, fluoroquinolones, aminoglycosides, macrolides, tetracycline, cephalosporins and sulfonamide. LS 25 and SS 3 show resistance against beta lactams and


glycopeptides again with the common mode of action, i.e. inhibition of cell wall synthesis. They are closely related members with respect to similarities in 16S rRNA gene sequences which justifies the observations of [48], that evolutionarily related bacteria have greater chance of being selected for in polluted environments either due to the presence of antibiotics or by the process of co selection of other pollutants (Fig. 4).

This study brings to light three important insights towards the ever-increasing burden of antimicrobial resistance in the environment, specially in wastewater. The first is the identification of multidrug resistant *E. coli* which has been included as one of the most important AMR indicators along with methicillin-resistant *Staphylococcus aureus* (MRSA) from 2019 onwards.


WHO reports (<https://www.who.int/news-room/fact-sheets/detail/antimicrobial-resistance>) suggests that 25 countries, territories and areas have provided data to the Global Antimicrobial Resistance and Use Surveillance System (GLASS) on blood-stream infections due to MRSA and 49 countries provided data on bloodstream infections due to *E. coli* where the median rate observed for methicillin-resistant *S. aureus* was 12.11% with an inter quartile range (IQR) of 6.4–26.4 and that for *E. coli* resistant to third generation cephalosporins was 36.0% (IQR -15.2–63.0). Thus the multidrug resistant isolate obtained from this study can also provide important insights towards the resistant gene profiles if studied further. The second important



**Fig. 2. Antibiotic resistant bacteria isolated from a rural setting in West Bengal, India (Graphical representation of the number of resistant isolates in LS sample set against various antibiotics; the number of resistant isolates is represented on y-axis and the antibiotics on x-axis) (a) and Antibiotic resistant bacteria isolated from an urban setting in West Bengal, India (Graphical representation of the number of resistant isolates in SS sample set against various antibiotics; the number of resistant isolates is represented on y-axis and the antibiotics on x-axis) (b)**



**Fig. 3. Phylogenetic analysis. Four distinct clades were identified in the phylogenetic tree (Clade 1 -LS 4, SS 7, LS 17, LS 9, SS 19; Clade 2- LS 6, LS 2, LS 21; Clade 3- LS 14, SS 20, LS 12; Clade 4- LS 23, SS 3 and LS 25**



**Fig. 4. Antibiotic resistance patterns of bacteria isolated from wastewaters collected from rural and urban areas of West Bengal, India. (Graphical representation of % antibiotic resistance on y-axis plotted against various classes of antibiotics on x-axis)**



**Table 2. Description of isolates represented in the phylogenetic tree, along with their sample IDs and GenBank Accession numbers**

Phylogenetic code	Sample ID	Accession number	Bacterial identity
Sample_1	LS 4	MW380613	<i>Shigella flexneri</i>
Sample_2	LS 6	MW380614	<i>Klebsiella pneumoniae</i>
Sample_3	LS 2	MW380615	<i>Escherichia coli</i>
Sample_4	LS 23	MW380616	<i>Lysinibacillus fusiformis</i>
Sample_5	LS 14	MW380617	<i>Bacillus australimaris</i>
Sample_6	SS 20	MW380618	<i>Bacillus pumilus</i>
Sample_7	LS 21	MW380619	<i>Comamonas aquatica</i>
Sample_8	LS 12	MW380620	<i>Bacillus safensis</i>
Sample_9	SS 3	MW380621	<i>Terribacillus halophilus</i>
Sample_10	SS 7	MW380622	<i>Citrobacter freundii</i>
Sample_11	LS 17	MW380623	<i>Escherichia coli</i>
Sample_12	LS 25	MW380624	<i>Oceanobacillus caeni</i>
Sample_13	LS 9	MW380625	<i>Escherichia coli</i>
Sample_14	SS 19	MW380626	<i>Comamonas aquatica</i>

isolate exhibiting resistance is *Lysinibacillus fusiformis* which have been reported in environmental samples but with very little resistant properties. Nonribosomal peptide synthetases (NRPS) and polyketide synthases (PKS) have been isolated from *Lysinibacillus fusiformis* both of which regulate the synthesis of antimicrobial compounds in the organism [49]. Thus the development of multidrug resistance in the isolate obtained in this study, probably indicates towards a mechanism in which the bacteria is able to survive competition in a particular wastewater niche as well as fortify its drug resistance mechanisms. The third interesting isolate is *Comamonas aquatica*, found in both urban and rural samples and resistant to beta lactams which has not been reported earlier. However, recent reports suggest the presence of antibiotic resistant *Comamonas testosteroni* in hospital set ups and in acute appendicitis [50]. It seems that the bacteria is increasing its resistance repertoire which is alarming in terms of the potential threat which it might pose in the coming years.

#### 4. CONCLUSION

The results obtained in our analyses, leads us to believe that untreated medical waste and anthropogenic waste disposal can lead to the propagation of different antibiotic resistant strains in wastewater environments both in urban and rural set ups. It is quite evident that the isolates from rural area showed high rates of resistance as compared to that of urban area. Excreted antibiotics can end up in wastewater treatment plants which are capable of degrading the

compound only partially. At the same time, such facilities probably serve as hotspots of horizontal gene transfer between bacterial species and even a few such strains if released in the environment can contribute negatively [51]. This reflects on the lack of awareness towards proper antibiotic usage in rural areas as compared to that in urban areas. This reflects on the need to practice efficient waste disposal and wastewater treatment policies. In natural settings, bacteria harboring antibiotic resistance genes are part of complex communities where they interact with other species [52].

This study should enable us to understand the route of spread of antibiotic resistant isolates in urban and rural wastewaters thus necessitating public awareness regarding measures of hygiene. Further, it should also enable us to formulate a monitoring program for evaluating wastewater health using a combination of culture independent and dependent molecular techniques to limit the spread of these emerging contaminants under tropical conditions.

#### SUPPLEMENTARY MATERIALS

Supplementary material is available in the following link: <https://www.journaljpri.com/index.php/JPRI/libraryFiles/downloadPublic/22>

#### CONSENT

It is not applicable.

#### ETHICAL APPROVAL

It is not applicable.

## ACKNOWLEDGEMENTS

The work is supported by the funds provided by the Department of Science and Technology and Biotechnology, Govt. of West Bengal, India project 236 (Sanc./ST/P&T/1G-39/2017). The authors hereby acknowledge the Department of Microbiology and Department of Biotechnology St. Xavier's College, Kolkata for providing the necessary facilities for conducting this study.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Kraemer SA, Ramachandran A, Perron GG. Antibiotic pollution in the environment: From microbial ecology to public policy. *Microorganisms*. 2019;7(6):180. DOI: 10.3390/microorganisms7060180.
- Klein EY, Tseng KK, Pant S, Laxminarayan R. Tracking global trends in the effectiveness of antibiotic therapy using the Drug Resistance Index. *BMJ Glob Health*. 2019;4(2):e001315. DOI:10.1136/bmjgh-2018-001315
- Taneja N, Sharma M. Antimicrobial resistance in the environment: The Indian scenario. *Indian J Med Res*. 2019;149(2):119-128. DOI:10.4103/ijmr.IJMR\_331\_18
- Manyi-Loh C, Mamphweli S, Meyer E, Okoh A. Antibiotic use in agriculture and its consequential resistance in environmental sources: Potential public health implications. *Molecules*. 2018;23(4):795. DOI: 10.3390/molecules23040795
- Von Wintersdorff CJ, Penders J, van Niekerk JM, Mills ND, Majumder S, van Alphen LB, et al. Dissemination of Antimicrobial Resistance in Microbial Ecosystems through Horizontal Gene Transfer. *Front Microbiol*. 2016;7:173. DOI:10.3389/fmicb.2016.00173
- Cycoń M, Mroziak A, Piotrowska-Seget Z. Antibiotics in the Soil Environment-Degradation and Their Impact on Microbial Activity and Diversity. *Front Microbiol*. 2019;10:338. DOI: 10.3389/fmicb.2019.00338.
- Chaudhry SB, Veve MP, Wagner JL. Cephalosporins: A Focus on Side Chains and  $\beta$ -Lactam Cross-Reactivity. *Pharmacy (Basel)*. 2019;7(3):103. DOI: 10.3390/pharmacy7030103.
- Cavicchioli R, Ripple WJ, Timmis KN, Azam F, Bakken LR, Baylis M, et al. Scientists' warning to humanity: microorganisms and climate change. *Nat Rev Microbiol*. 2019;17(9):569-586. DOI:10.1038/s41579-019-0222-5.
- Bengtsson-Palme J, Kristiansson E, Larsson DGJ. Environmental factors influencing the development and spread of antibiotic resistance. *FEMS Microbiol Rev*. 2018;42(1):fux053. DOI: 10.1093/femsre/fux053.
- Exner M, Bhattacharya S, Christiansen B, Gebel J, Goroncy-Bermes P, Hartemann P, et al. Antibiotic resistance: What is so special about multidrug-resistant Gram-negative bacteria?. *GMS Hyg Infect Control*. 2017;12:Doc05. DOI: 10.3205/dgkh000290.
- Karkman A, Do TT, Walsh F, Virta MPJ. Antibiotic-Resistance Genes in Waste Water. *Trends Microbiol*. 2018;26(3):220-228. DOI: 10.1016/j.tim.2017.09.005.
- Grennan D, Varughese C, Moore NM. Medications for Treating Infection. *JAMA*. 2020;323(1):100. DOI: 10.1001/jama.2019.17387.
- Rizzo L, Manaia C, Merlin C, Schwartz T, Dagot C, Ploy MC, et al. Urban wastewater treatment plants as hotspots for antibiotic resistant bacteria and genes spread into the environment: a review. *Sci Total Environ*. 2013;447:345-360. DOI: 10.1016/j.scitotenv.2013.01.032.
- The center for disease, dynamics economics & policy. Resistance map: Antibioticresistance;2020. Available: <https://resistancemap.cddep.org/AntibioticResistance.php>. Date accessed: January 15, 2021
- Langford BJ, So M, Raybardhan S, Leung V, Westwood D, MacFadden DR, et al. Bacterial co-infection and secondary infection in patients with COVID-19: a living rapid review and meta-analysis. *Clin Microbiol Infect*. 2020;(12):1622-1629. DOI: 10.1016/j.cmi.2020.07.016.
- Singh M, Ganguli S, Ghosh MM. Comparative metagenomic dataset of hospital effluent microbiome from rural and urban hospitals in West Bengal. *Data Brief*. 2019;25:104264. DOI: 10.1016/j.dib.2019.104264.
- Available: [http://kmdaonline.org/home/aar\\_2011](http://kmdaonline.org/home/aar_2011)

18. Lokhande RS, Singare PU, Pimple DS. Study on Physico-chemical Parameters of Waste Water Effluents from Talaja Industrial Area of Mumbai, India. *Int. J. Ecosyst.* 2011;1(1):1-9. DOI:10.5923/j.ije.20110101.01
19. Bergey DH, Holt JG. *Bergey's manual of determinative bacteriology*. 9th ed. Baltimore: Williams & Wilkins; 1994.
20. Gerhardt P, Murray RGE, Wood WA, Krieg NR. *Methods for general and molecular bacteriology*. Washington D.C: American Society for Microbiology; 1994.
21. Clinical and Laboratory Standards Institute (CLSI). *Performance Standards for Antimicrobial Susceptibility Testing*. 30th ed. CLSI supplement M100 (ISBN 978-1-68440-066-9 [Print]; ISBN 978-1-68440-067-6 [Electronic]). Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087 USA, 2020.
22. Available: <https://www.ncbi.nlm.nih.gov/>
23. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol.* 2011;7:539. DOI: 10.1038/msb.2011.75.
24. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 2004;32(5):1792-1797. Available:<https://doi.org/10.1093/nar/gkh340>.
25. Felsenstein J. Phylogenies and the Comparative Method. *Am. Nat.* 1985;125(1):1-15.
26. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol.* 1987;4(4):406-25. DOI: 10.1093/oxfordjournals.molbev.a040454.
27. Felsenstein J. Evolutionary trees from DNA sequences: A maximum likelihood approach. *J Mol Evol.* 1981;17(6):368-376.
28. Nguyen CC, Hugie CN, Kile ML. Association between heavy metals and antibiotic-resistant human pathogens in environmental reservoirs: A review. *Front Environ Sci Eng.* 2019;13:46. DOI:10.1007/s11783-019-1129-0.
29. Santos-Lopez A, Marshall CW, Scribner MR, Snyder DJ, Cooper VS. Evolutionary pathways to antibiotic resistance are dependent upon environmental structure and bacterial lifestyle. *Elife.* 2019;8:e47612. DOI: 10.7554/eLife.47612.
30. Estrada-García T, Cerna JF, Paheco-Gil L, Velázquez RF, Ochoa TJ, Torres J, et al. Drug-resistant diarrheogenic *Escherichia coli*, Mexico. *Emerg Infect Dis.* 2005;11(8):1306-8. DOI: 10.3201/eid1108.050192.
31. Sabir S, Anjum AA, Ijaz T, Ali MA, Khan MUR, Nawaz M. Isolation and antibiotic susceptibility of *E. coli* from urinary tract infections in a tertiary care hospital. *Pak J Med Sci.* 2014;30(2):389-92.
32. Caneiras C, Lito L, Melo-Cristino J, Duarte A. Community- and Hospital-Acquired *Klebsiella pneumoniae* Urinary Tract Infections in Portugal: Virulence and Antibiotic Resistance. *Microorganisms.* 2019;7(5):138. DOI: 10.3390/microorganisms7050138.
33. Manikandan C, Amsath A. Antibiotic susceptibility pattern of *Klebsiella pneumoniae* isolated from urine samples. *Int. J. Curr. Microbiol. App. Sci.* 2013;2(8):330-337.
34. Ranjbar R, Farahani A. *Shigella*: Antibiotic-Resistance Mechanisms And New Horizons For Treatment. *Infect Drug Resist.* 2019;12:3137-3167. DOI: 10.2147/IDR.S219755.
35. Ahmed I, Yokota A, Yamazoe A, Fujiwara T. Proposal of *Lysinibacillus boronitolerans* gen. nov. sp. nov., and transfer of *Bacillus fusiformis* to *Lysinibacillus fusiformis* comb. nov. and *Bacillus sphaericus* to *Lysinibacillus sphaericus* comb. nov. *Int J Syst Evol Microbiol.* 2007;57(Pt 5):1117-1125. DOI: 10.1099/ijs.0.63867-0.
36. Priest FG, Goodfellow M, Todd C. A numerical classification of the genus *Bacillus*. *J Gen Microbiol.* 1988;134(7):1847-82. DOI: 10.1099/00221287-134-7-1847. PMID: 3246588.
37. Shivaji S, Chaturvedi P, Suresh K, Reddy GSN, Dutt CBS, Wainwright M, et al. *Bacillus aerius* sp. nov., *Bacillus aerophilus* sp. nov., *Bacillus stratosphericus* sp. nov. and *Bacillus altitudinis* sp. nov., isolated from cryogenic tubes used for collecting air samples from high altitudes. *Int J Syst Evol Microbiol.* 2006;56(Pt 7):1465-1473. DOI: 10.1099/ijs.0.64029-0.
38. Liu Y, Lai Q, Dong C, Sun F, Wang L, Li G, et al. Phylogenetic diversity of the *Bacillus*

- pumilus* group and the marine ecotype revealed by multilocus sequence analysis. PLoS One. 2013;8(11):e80097.
39. Branquinho R, Klein G, Kämpfer P, Peixe LV. The status of the species *Bacillus aerophilus* and *Bacillus stratosphericus*. Request for an Opinion. Int J Syst Evol Microbiol. 2015;65(Pt 3):1101. DOI: 10.1099/ijs.0.000004.
  40. Farooq S, Farooq R, Nahvi N. *Comamonas testosteroni*: Is It Still a Rare Human Pathogen? Case Rep Gastroenterol. 2017;11(1):42-47. DOI: 10.1159/000452197.
  41. Steinberg JP, Burd EM. Other Gram-Negative and Gram-Variable Bacilli. In: Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases. 8th ed; 2015.
  42. Nam JH, Bae W, Lee DH. *Oceanobacillus caeni* sp. nov., isolated from a *Bacillus*-dominated wastewater treatment system in Korea. Int J Syst Evol Microbiol. 2008;58(Pt 5):1109-1113. DOI: 10.1099/ijs.0.65335-0.
  43. Liu L, Chen D, Liu L, Lan R, Hao S, Jin W, et al. Genetic Diversity, Multidrug Resistance, and Virulence of *Citrobacter freundii* From Diarrheal Patients and Healthy Individuals. Front Cell Infect Microbiol. 2018;8:233. DOI: 10.3389/fcimb.2018.00233.
  44. An SY, Asahara M, Goto K, Kasai H, Yokota A. *Terribacillus saccharophilus* gen. nov., sp. nov. and *Terribacillus halophilus* sp. nov., spore-forming bacteria isolated from field soil in Japan. Int J Syst Evol Microbiol. 2007;57(Pt 1):51-55. DOI: 10.1099/ijs.0.64340-0.
  45. Conte D, Palmeiro JK, da Silva Nogueira K, de Lima TM, Cardoso MA, Pontarolo R, et al. Characterization of CTX-M enzymes, quinolone resistance determinants, and antimicrobial residues from hospital sewage, wastewater treatment plant, and river water. Ecotoxicol Environ Saf. 2017;136:62-69. DOI: 10.1016/j.ecoenv.2016.10.031.
  46. Lupo A, Coyne S, Berendonk TU. Origin and evolution of antibiotic resistance: the common mechanisms of emergence and spread in water bodies. Front Microbiol. 2012;3:18. DOI: 10.3389/fmicb.2012.00018.
  47. Gullberg E, Cao S, Berg OG, Ilbäck C, Sandegren L, Hughes D, et al. Selection of resistant bacteria at very low antibiotic concentrations. PLoS Pathog. 2011;7(7):e1002158. DOI: 10.1371/journal.ppat.1002158.
  48. Martinez JL, Sanchez MB, Martinez-Solano L, Hernandez A, Garmendia L, Fajardo A, et al. Functional role of bacterial multidrug efflux pumps in microbial natural ecosystems. FEMS Microbiol Rev. 2009;33:430-449. DOI: 10.1111/j.1574-6976.2008.00157.x.
  49. Tartar SA, Tartar T. A Rare Pathogen in Acute Appendicitis: Two Cases with *Comamonas testosteroni* Infection and Literature Review. J Pediatr Infect Dis. 2020;15(02):110-112. DOI: 10.1055/s-0038-1641604
  50. Zothanpuia, Passari AK, Gupta VK, Singh BP. Detection of antibiotic-resistant bacteria endowed with antimicrobial activity from a freshwater lake and their phylogenetic affiliation. PeerJ. 2016;4:e2103. DOI: 10.7717/peerj.2103.
  51. Hiltunen T, Virta M, Laine AL. Antibiotic resistance in the wild: an eco-evolutionary perspective. Philos Trans R Soc Lond B Biol Sci. 2017;372(1712):20160039. DOI: 10.1098/rstb.2016.0039.
  52. Munita JM, Arias CA. Mechanisms of Antibiotic Resistance. Microbiol Spectr. 2016;4(2):10.1128/microbiolspec.VMBF-0016-2015. DOI: 10.1128/microbiolspec.VMBF-0016-2015.

© 2021 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/77759>