



Response of Different Levels of Plant Growth Regulators on Gerbera (*Gerbera jamesonii*) cv. Pink Elegance under Protected Cultivation

Shinnu Sabu ^{a++*}, Samir E Topno ^b, Vijay Bahadur ^b,
Annjoe V Joseph ^b and Anupriya Paul ^c

^a Department of Floriculture & Landscaping, SHUATS, India.

^b Department of Horticulture, SHUATS, India.

^c Department of Mathematics & Statistics, SHUATS, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i7982>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118200>

Original Research Article

Received: 24/03/2024

Accepted: 29/05/2024

Published: 07/06/2024

ABSTRACT

An experiment was conducted to evaluate the influence of plant growth regulators on the growth, flower quality and yield of gerbera (*Gerbera jamesonii* Adlam) cv. Pink Elegance under a naturally ventilated polyhouse in the Department of Horticulture at SHUATS, Allahabad, Uttar Pradesh, India, during 2023-24. The design for experiment is Randomized Block Design with three replications and ten treatments. Gerbera plants were sprayed with GA₃, Cycocel and NAA at different concentrations of 50, 100 and 150 ppm along with control. The results from the study concluded that the treatment

⁺⁺ M.Sc. Scholar;

*Corresponding author: E-mail: shinnusabu123@gmail.com;

Cite as: Sabu, Shinnu, Samir E Topno, Vijay Bahadur, Annjoe V Joseph, and Anupriya Paul. 2024. "Response of Different Levels of Plant Growth Regulators on Gerbera (*Gerbera Jamesonii*) Cv. Pink Elegance under Protected Cultivation". *Journal of Advances in Biology & Biotechnology* 27 (7):220-26. <https://doi.org/10.9734/jabb/2024/v27i7982>.

GA₃ @150 ppm was found in superior among other treatments in terms of plant height, number of leaves per plant, plant spread, leaf area, leaf area index, number of flowers per plant, flower weight, flower diameter, root length, stalk length, shelf life, flower yield per m², flower yield per 1000 polybags and minimum days to first flowering. Thus, the 150 ppm GA₃ treatment was identified as the most effective for promoting both excellent vegetative and reproductive growth. Thus, GA₃ @150 ppm can be recommended for plant growth and flowering of Gerbera proving the multifaceted role of plant growth regulator (PGR) in gerbera cultivation, helping growers optimize plant growth, flowering and yield while maintaining high-quality flower production.

Keywords: Growth regulators; cycocel; gibberellic acid (GA₃); gerbera; growth; naphthelen acetic acid (NAA); yield.

1. INTRODUCTION

“Gerbera (*Gerbera jamesonii* Adlam), a prominent cut flower, ranks fifth among the top ten cut flowers in the global market and is extensively cultivated under protected conditions in regions like Bangalore, Pune, and Uttarakhand, India. Gerbera is native to tropical Asia and Africa, this species belongs to the Asteraceae family and known by other various common names such as Transvaal Daisy, Barberton Daisy, or African Daisy and can be grown under a wide range of climatic conditions” [1]. “Discovered by Gronovious and named in honor of German naturalist Traugott Gerber, gerbera is valued for its wide range of colours, including yellow, red, orange, cream, white, and various shades in between” [2]. These features, along with its use in floral arrangements and decorations, make it a popular choice in India [3,4,5].

“Gerbera is a dwarf herbaceous perennial, characterized by solitary flower heads on long slender stalks, which thrive in polyhouses and shade houses” [6]. “These environments enhance plant growth, producing larger, greener leaves with high dry matter content. The commercial production of gerbera is in high demand both nationally and internationally, especially during winter when few other flowers bloom” [7].

“Gerbera flowers are in high demand nationwide due to their enduring nature, elegant appearance, resilience, and export potential. Optimizing the application of plant growth regulators (PGRs) can lead to increased productivity and improved quality of gerbera flowers, thus enhancing the economic viability of gerbera production. Understanding the specific effects of different PGRs on gerbera growth, flowering, and fruiting can enable growers to tailor their application methods to maximize

desired outcomes while minimizing costs and environmental impacts” [8]. Moreover, research in this area can contribute to the development of sustainable cultivation practices by reducing the reliance on conventional chemical inputs and promoting environmentally friendly alternatives and gerbera plants can inform breeding efforts aimed at developing new cultivars with enhanced responsiveness to PGRs or intrinsic resistance to environmental stresses.

“Plant growth regulators have revolutionized the floriculture industry by significantly enhancing the growth, yield, and quality of ornamental crops like gerbera. PGRs, including gibberellic acid (GA₃), cycocel and naphthalene acetic acid (NAA), are chemical substances used in small amounts to regulate plant development and stimulate desired growth responses [9]. They help overcome growth limitations, stimulate flowering, and extend the vase life of flowers” [10]. Given the significance of PGRs, the present study aimed to determine the optimal concentrations of GA₃, cycocel and NAA for improving the growth, yield, and quality of gerbera under naturally ventilated polyhouses. This research is conducted to study the effect of different levels of PGR on growth, flowering and yield of Gerbera.

2. MATERIALS AND METHODS

The experiment was conducted during the winter season of 2023-24 at the Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Allahabad, Uttar Pradesh. The experimental site is located at a latitude of 20°15' North and a longitude of 60° East, at an altitude of 98 meters above mean sea level (MSL). The soil at the experimental site was sandy loam in texture with a pH of 6.87, an electrical conductivity (E.C) of 0.15 dS/m, and a carbon content of 43%. The available nitrogen,

phosphorus, and potassium contents were 216.62 kg/ha, 37.34 kg/ha, and 215.86 kg/ha, respectively. The experiment was laid out in a Randomized Block Design with three replications and ten treatments. Seedlings of gerbera cv. Pink Elegance, each with 4-5 leaves, were planted on raised beds on November 13, 2023 in polybags. Three plant growth regulators, gibberellic acid (GA₃), cycocel and naphthalene acetic acid (NAA), were used at three concentrations 50, 100 and 150ppm along with the control. The details of the treatment combinations are as follows: T₀- Control, T₁- GA₃ 50 ppm, T₂- GA₃ 100 ppm, T₃- GA₃ 150 ppm, T₄- Cycocel 50 ppm, T₅- Cycocel 100 ppm, T₆- Cycocel 150 ppm, T₇- NAA 50 ppm, T₈- NAA 100 ppm and T₉- NAA 150 ppm.

3. RESULTS AND DISCUSSION

3.1 Influence of Plant Growth Regulators on Growth Parameters

The data on growth parameters presented in Table 1 show that the maximum plant height (35.68 cm), number of leaves (30.96), plant spread (51.21 cm), leaf area (178.64m²) and leaf area index (3.77) were observed in treatment T₃ (GA₃ @ 150 ppm). In contrast, the minimum plant height (21.12 cm), number of leaves (18.16), plant spread (38.02 cm), leaf area (129.65 m²) was at par with T₄ (124.06m²) and leaf area index (1.71) were recorded in treatment T₀ (control). The significantly greater plant height, number of leaves, plant spread, leaf area and leaf area index in T₃ can be attributed to the effect of gibberellic acid, which promotes cell division and cell elongation, leading to enhanced vegetative growth. Similar findings regarding maximum plant height were reported by Nair et al. [11] and Dalal et al. [12] in gerbera. The maximum plant spread at 150 ppm was observed by Patra et al. [13] in gerbera. Sharma et al. [14] suggested that GA₃ at 100 ppm is highly effective in promoting vegetative growth in gladiolus and Dogra et al. [15] reported an increase in leaf area index due to GA₃ treatments. Similarly, Porwal et al. [16] observed the impact of plant growth regulators on Damask Rose, noting that GA₃ at 250 ppm resulted in the maximum plant height, number of shoots per plant, and plant spread. Additionally, GA₃ promotes the increase in the photosynthetic area by inducing cell elongation and hyponasty of leaves, which affects their orientation and ultimately contributes to an increase in leaf area [17].

3.2 Influence of Plant Growth Regulators on Flowering Parameters

The data on qualitative parameters presented in Table 2 indicate that the minimum number of days to first flower bud emergence or earliness, which occurred in 69.33 days, was observed in T₃ (GA₃ @ 150 ppm), while T₀ (control) took the maximum number of days (93.11days) to produce a visible flower bud. The maximum number of flowers per plant (12.00), flower weight (40.12), flower diameter (12.31), stalk length (47.88), shelf life (9.95), and root length (26.80) was observed with T₃ - GA₃ @150 ppm. Conversely, the minimum number of flowers per plant (4.67), flower weight (26.72), flower diameter (7.24), stalk length (35.55), shelf life (5.63) and root length (14.86) was found in treatment T₀ (control).

Gibberellic acid is a plant growth regulator known to play a crucial role in various physiological processes, including cell division, elongation, and differentiation. When applied exogenously, GA₃ can stimulate these processes, leading to significant changes in plant morphology and development [18]. In the case of gerbera plants, the application of GA₃ at 150 ppm stimulated the cell expansion and enlargement of floral structures which have accelerated the initiation and emergence of flower buds, resulting in earlier flowering, promoted flower bud formation, enhanced floral organ development, increased flower weight and diameter [19]. Moreover, GA₃ (150ppm) treated plants displayed enhanced root and stalk growth, with longer root systems and taller stalks, which can support larger and more abundant flowers. It also improved the post-harvest characteristics such as shelf life, as GA₃ at 150ppm treated flowers exhibited delayed senescence and prolonged freshness compared to untreated flowers [20]. Overall, the application of GA₃ at 150 ppm effectively optimized qualitative parameters in gerbera plants, resulting in enhanced flower quality and overall plant performance.

Patel et al. [21] observed similar results in gerbera, with GA₃ at 150 ppm resulting in the fewest days to first flower initiation (108.33 days), maximum flower diameter (8.76 cm), flower weight (5.93 g), and shelf life of flowers (8.00 days). GA₃ also increased the number of suckers, and foliar application of 150 ppm GA₃ was found to be the best for enhancing plant growth, yielding the maximum number of cut blooms with longer stalks and larger flower sizes,

as reported by Sharifuzzaman et al. [22] in chrysanthemum, Jamal et al. [23] in gerbera and Sainath et al. [24] in chrysanthemum observed similar earliness in the full opening of flowers due to GA₃ treatments. Chauhan et al. [25] found that GA₃ at 150 ppm led to the earliest appearance of the first flower bud (50.98 days) and the largest

flower diameter (11.37 cm) in gerbera. Furthermore, Salem et al. [26] reported that in gerbera, the highest yield and quality parameters (number of flowers per plant, stalk length, flower bud diameter, and stalk diameter), as well as the shortest time to flower bud emergence and first flowering, were observed with GA₃ at 150 ppm.

Table 1. Effect of plant growth regulators on growth parameters of gerbera (*Gerbera jamesonii* cv. Pink elegance)

Treatments	Plant height (cm)	Number of leaves/plant	Plant spread (cm)	Leaf Area (m ²)	Leaf Area Index (cm ²)
T ₀	21.12	18.16	38.02	129.65	1.71
T ₁	31.66	24.86	47.02	146.40	2.78
T ₂	33.30	28.79	49.68	171.19	3.19
T ₃	35.68	30.96	51.21	178.64	3.77
T ₄	27.43	19.25	39.72	124.06	2.04
T ₅	29.77	21.67	41.35	152.34	2.46
T ₆	29.33	23.56	44.85	164.18	2.53
T ₇	24.04	22.24	42.02	164.48	2.36
T ₈	25.49	25.91	43.35	153.13	2.94
T ₉	27.04	26.73	45.35	146.58	3.12
F-Test	S	S	S	S	S
S.Ed (+)	0.94	1.06	0.87	2.69	0.06
CD_(0.05)	2.00	2.24	1.84	5.70	0.13

Note: T₀= control; T₁= GA₃ 50 ppm; T₂= GA₃ 100 ppm; T₃= GA₃ 150 ppm. T₄= cycocel 50 ppm; T₅= cycocel 100 ppm; T₆= cycocel 150 ppm. T₇= NAA 50 ppm; T₈= NAA 100 ppm; T₉= NAA 150 ppm

Table 2. Effect of plant growth regulators on floral parameters of gerbera (*Gerbera jamesonii* cv. Pink elegance)

Treatments	Days to first flowering	Flower weight (g)	Flower diameter (cm)	Stalk length (cm)	Shelf life (days)	Root length (cm)
T ₀	93.11	26.72	7.24	35.55	5.63	14.86
T ₁	72.48	36.99	10.65	44.35	7.67	23.66
T ₂	70.27	38.64	11.72	45.93	8.69	24.85
T ₃	69.33	40.12	12.31	47.88	9.95	26.80
T ₄	88.01	29.39	7.83	36.22	6.63	17.42
T ₅	84.88	31.03	8.09	38.47	6.61	18.19
T ₆	80.29	32.24	8.75	38.69	6.34	18.36
T ₇	77.73	34.72	9.98	39.33	7.52	19.44
T ₈	75.26	34.81	10.26	40.62	8.31	22.21
T ₉	78.33	35.36	10.31	42.31	8.66	22.30
F-Test	S	S	S	S	S	S
S.Ed (+)	1.78	0.64	0.21	1.13	0.44	0.52
CD_(0.05)	3.77	1.35	0.45	2.40	0.94	1.11

Note: T₀= control; T₁= GA₃ 50 ppm; T₂= GA₃ 100 ppm; T₃= GA₃ 150 ppm. T₄= cycocel 50 ppm; T₅= cycocel 100 ppm; T₆= cycocel 150 ppm. T₇= NAA 50 ppm; T₈= NAA 100 ppm; T₉= NAA 150 ppm

Table 3. Effect of plant growth regulators on yield parameters of gerbera (*Gerbera jamesonii* cv. Pink elegance)

Treatments	Flowers number per plant	Flower yield per m ²	Flower yield per 1000 polybags
T ₁₀	4.67	65.15	4670
T ₁	8.23	99.39	8230
T ₂	10.00	107.58	10000
T ₃	12.00	112.89	12000
T ₄	6.86	77.58	6860
T ₅	7.00	76.11	7000
T ₆	7.57	90.23	7570
T ₇	7.61	82.05	7610
T ₈	8.12	86.54	8120
T ₉	9.20	92.15	9200
F-Test	S	S	S
S.Ed (±)	0.70	1.55	180.414
CD (5%)	1.48	3.28	381.972

Note: T₀= control; T₁= GA₃ 50 ppm; T₂= GA₃ 100 ppm; T₃= GA₃ 150 ppm. T₄= cycocel 50 ppm; T₅= cycocel 100 ppm; T₆= cycocel 150 ppm. T₇= NAA 50 ppm; T₈= NAA 100 ppm; T₉= NAA 150 ppm

3.3 Influence of Plant Growth Regulators on Yield Parameters

The data on yield parameters presented in Table 3 indicate that the maximum flower yield per m² (112.89) and maximum flower yield per 1000 polybags (12000) was observed with T₃ - GA₃ @150 ppm. While, the minimum flower yield per m² (65.15) and flower yield per 1000 polybags (4670) were recorded in treatment T₀ (control). The increase in flower yield could be attributed to the fact that plants treated with gibberellic acid produced a greater number of leaves and number of flowers. This increased leaf production likely led to the production and accumulation of more photosynthates, which were then redirected to the floral organs, resulting in a higher number of flowers and flower weight. These findings are consistent with the results reported by Dalal et al. [12] in gerbera, Tariq et al. [27] in gladiolus, Patel et al. [28] in African marigold, Patel et al. (2016) in chrysanthemum, Kumar et al. [29] and Sainath et al. [24] in chrysanthemum [30,31].

4. CONCLUSION

From the present investigation, it is concluded that among the different treatments, the treatment T₃ - GA₃ @150 ppm was found in superior among other treatments in terms of Plant height (33.30 cm), Number of leaves per plant (30.96), plant spread (49.68), leaf area (152.34m²), leaf area index (3.77), number of flowers per plant (12.00), flower weight (40.12g), flower diameter (12.31cm), root length (26.80cm), stalk length (47.88cm), shelf life

(9.95), flower yield per 1000 polybags (12000) and minimum days to first flowering (69.33). Thus, GA₃ @150 ppm can be recommended for plant growth and flowering of Gerbera proving the multifaceted role of plant growth regulator in gerbera cultivation, helping growers optimize plant growth, flowering and yield while maintaining high-quality flower production

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Swarup V. Garden Flowers, National Book Trust. 1997;184-185.
2. Prasad KV. Bulbous ornamental crops: Status of global trade. Floriculture Today. 2007;11(8): 22-36.
3. Rashid Z, Ahmad Ahngar T, Sabiya B, Sabina N, Khuroo NS, Shabeena M, Rakshanda AA. Bhat R, Rasool F-U-, Shafeeq H, Dar ZA, Jan S. Influence of plant growth regulators and micronutrients on growth, yield and quality of sorghum under temperate conditions. J. Exp. Agric. Int. 2021 Oct. 11 [cited 2024 May 28];43(9):18-23. Available:https://journaljeai.com/index.php/JEAI/article/view/1878
4. Vishal, Bahadur V, Manjeet. Impact of plant growth regulators on growth, yield and quality of strawberry (*Fragaria × ananassa* Duch.) cv. Winter Dawn. Int. J.

- Plant Soil Sci. 2023 Aug. 14 [cited 2024 May 28];35(19):31-6.
Available: <https://journalijpss.com/index.php/IJPSS/article/view/3520>
5. Khan N, Bano A, Shahid MA, Nasim W, Ali Babar MD. Interaction between PGPR and PGR for water conservation and plant growth attributes under drought condition. *Biologia*. 2018 Nov;73:1083-98.
 6. Sahu MK, Tirkey T, Sharma G, Tiwari A, Kushram T. Effect of foliar application of micronutrients on growth and flower production of gerbera under protected condition. *Journal of Pure and Applied Microbiology*. 2016;10(4):3099-3103.
 7. Darras A. Overview of the dynamic role of specialty cut flowers in the international cut flower market. *Horticulturae*. 2021;7(3):51.
 8. Fawzy ZF, El-Shal ZS, Yunsheng L, Zhu O, Sawan OM. Response of gerbera plants to foliar spraying of some bio-stimulants under sandy soil condition. *Applied Scientific Research*. 2012;8(2):770-776.
 9. Wu X, Gong D, Zhao K, Chen D, Dong Y, Gao Y, Hao GF. Research and development trends in plant growth regulators. *Advanced Agrochem*. 2024;3(1):99-106.
 10. Thakur R, Thakur R, Chandermohan NC, Kanwar B. Effect of gibberellic acid and benzyl adenine on ornamental plants, *The Pharma Innovation Journal*. 2023;12(2):101-105.
 11. Nair SA, Singh V, Sharma TVRS. Effect of plant growth regulators on yield and quality of gerbera under Bay Island conditions. *Indian Journal of Horticulture*. 2002;59(1):100-105.
 12. Dalal SR, Somavanshi AV, Karale GD. Effect of gibberellic acid on growth, flowering, yield and quality of Gerbera under polyhouse condition; *International journal of Agriculture Sciences*. 2009;5(2):355-356.
 13. Patra SK, Beura S, Shasani T. Efficacy of GA₃ on growth and flowering regulation of *In vitro* raised hybrid Gerbera under shade net. *Agric. Sci. Digest*. 2015;35(3):173-177.
 14. Sharma JR, Gupta RB, Panwar RD. Growth, flowering and corm production of gladiolus cv. Friendship as influenced by foliar application of nutrients and growth regulators. *Journal of Ornamental Horticulture*. 2004;7(3-4):154–158.
 15. Dogra S, Pandey RK, Bhat DJ. Influence of gibberellic acid and plant geometry on growth, flowering production in gerbera under Jammu agroclimate, *International Journal of Pharma and Bio Sciences*. 2012;3(4):1083-1090.
 16. Porwal R, Nagda CL, Pundir JPS. Influence of plant growth regulator on vegetative growth and flower earliness of Damask rose. *South Indian Hort*. 2002;50(1-3):119-123.
 17. Dutta SC. *Plant physiology*. Wiley Eastern Limited. New Age International Limited. 2015;466-558.
 18. Ridha AAS, Saravanan S, Prasad VM. Effect of gibberellic acid spraying on yield and flowers of gerbera (*Gerbera jamesonii*) Cv. Dennis. *International J. of Scientific Res*. 2016;5(3):234-236.
 19. Saini S, Arora S. Effect of gibberellic acid on growth, flowering, yield and quality of gerbera under polyhouse conditions, *Indian Nutrient Digest*. 2016;11:233-236.
 20. Zosser CN, Sangma Devi S, Urfi F. Effect of plant growth regulators on growth, yield and flower quality of Gerbera (*Gerbera jamesonii* L.) cv. pink elegance under naturally ventilated polyhouse (NVPH), *International Journal of Current Microbiology and Applied Sciences*. 2017;6(10):468-476.
 21. Patel BB, Desai JR, Patel GD, Patel HF. Influence of foliar application of nitrogen and plant growth regulators on growth, flowering production of gerbera, *Bioinfolet. Journal*. 2013;10(2A):415 – 417.
 22. Sharifuzzaman SM, Ara KA, Rahman MH, Kabir K, Talukdar MB. Effect of GA₃, CCC and MH on vegetative growth, flower yield and quality of Chrysanthemum. *Int. J. Expt. Agric*. 2011;2(1):17-20.
 23. Jamal Uddin AFM, Mehraj H, Taufique T, Ona AF, Parvin S. Foliar application of gibberellic Acid on growth and flowering of gerbera cultivars, *Journal of Bioscience and Agriculture Research*. 2014;02(01):52-58.
 24. Sainath DS, Uppar VS, Patil VK, Deshpande, Husije R. Effect of different growth regulators on seed yield and quality attributes in annual chrysanthemum (*Chrysanthemum coronarium* L). *Journal of Agricultural Sciences*. 2018;27(2):131-134.
 25. Chauhan RV, Kav KP, Babariya VJ, Pansuria PB, Savaliya AB. Effect of gibberellic acid on flowering and cut flower yield in Gerbera under protected condition.

- Asian Journal of Horticulture. 2014;9(2): 404-407.
26. Salem RAA, Saravanan S, Prasad VM. Effect of gibberellic acid spraying on yield and flowers of Gerbera (*Gerbera jamesonii*) cv. Dennis. International Journal of Scientific Research. 2016;5(3):2277-2280.
27. Tariq S, Hassan I, Akhtar N, Abbasi Jilani G. Effect of gibberellic acid on the vase life and oxidative activities in senescing cut gladiolus flowers. Article in Plant Growth Regulation. 2013;72:89-95.
28. Patel S, Das AK, Dash DK. Effect of plant growth regulators on growth and yield attributes of african marigold (*Tagetes erecta* L.) International Education & Res. J. 2015;2(6):44-45.
29. Kumar M, Singh AK, Kumar A. Effect of plant growth regulators on flowering and yield attributes of African marigold (*Tagetes erecta* L.) cv. Pusanarangigainda, Plant Archives. 2014; 14(1):363-365.
30. Palei S, Das AK, Dash DK. Effect of plant growth regulators on flowering and yield attributes of African marigold (*Tagetes erecta* L.), International Education and Research Journal. 2016;2(6):2454-9916.
31. Patil D, Chopde N, Lokhande S, Bhande MH. Studies on response of African marigold to plant growth regulators for seed production. Plant Archives. 2016;16(1):423425.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/118200>