



Assessment of Genetic Variability, Heritability and Genetic Advance in White Onion (*Allium cepa* L.) for Growth, Bulb Yield, Quality and its Component Characteristics

Sunil Chavan ^{a+++*}, Hiremath S M ^{a#}, Gangadhara Doggalli ^{b†},
Vinutha Patil S ^{c†}, Ajay S M ^{a†}, Mahesh A Vernekar ^{a†},
Kavya ^{d‡} and Vinodh Kumar P N ^{e†}

^a Department of Horticulture, UAS, Dharwad-580005, India.

^b Department of Genetics and Plant Breeding, University of Agricultural Sciences, Dharwad, 580005, India.

^c Department of Genetics and Plant Breeding, University of Agricultural Sciences, Raichur, India.

^d ICAR –IGFRI, SRRS, University of Agricultural Sciences, Dharwad, 580005, India.

^e Department of Genetics and Plant Breeding, ICAR- IARI, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i91389>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/121982>

Original Research Article

Received: 25/06/2024

Accepted: 27/08/2024

Published: 06/09/2024

⁺⁺Msc. Scholar;

[#]Professor;

[†]PhD Scholar;

[‡]Project Assistant;

^{*}Corresponding author: E-mail: chavandani143@gmail.com;

Cite as: Chavan, Sunil, Hiremath S M, Gangadhara Doggalli, Vinutha Patil S, Ajay S M, Mahesh A Vernekar, Kavya, and Vinodh Kumar P N. 2024. "Assessment of Genetic Variability, Heritability and Genetic Advance in White Onion (*Allium Cepa* L.) for Growth, Bulb Yield, Quality and Its Component Characteristics". *Journal of Advances in Biology & Biotechnology* 27 (9):1194-99. <https://doi.org/10.9734/jabb/2024/v27i91389>.

ABSTRACT

The present investigation was carried out at Main Agricultural Research Station, University of Agricultural Sciences, Dharwad, Karnataka, during the latekharif season of 2023-2024. The study was undertaken to evaluate the variability present among the latekharif genotypes and predict the results for hybridization programme. Twenty-six genotypes of onion were used for the experiment. Results indicated substantial phenotypic variability across the genotypes, with traits such as marketable yield (t/ha) showing high genotypic and phenotypic coefficient of variation (GCV and PCV). Broad-sense heritability estimates ranged widely, emphasizing genetic control for traits like polar diameter (cm) and marketable yield (t/ha). Genetic advance, as a percentage of mean, highlighted significant annual gains in fresh weight of plant (g) and average weight of bulb (g). The findings underscore the potential for onion improvement through selective breeding strategies targeting traits with high heritability and genetic advance. This research contributes valuable insights into enhancing onion productivity and quality traits crucial for sustainable agricultural practices and global food security.

Keywords: Onion; variability; heritability; genetic advance.

1. INTRODUCTION

“Onion (*Allium cepa*L.) is one of the important culinary vegetable belongs to family *Alliaceae*, having chromosome number $2n=16$. It is a native of South West-Asia, from where it spread all over the world. The crop is mainly grown for local consumption and for export purposes. In the world, onion being the third most valuable vegetable after potato and tomato, and it is grown in more than 140 countries by covering an area of 54.8 lakh hectares with production of 1045.54 lakh tons and productivity of 23.06 t/ha” [1,2]. “Among the cultivated *Alliums* in India onion is a prominent export-oriented vegetable and forms the world's second largest producer after China. In India, it is being cultivated in an area of 1.43 million hectares, producing 26.09 million tonnes with a productivity of 18.23 tonnes per hectare” [3]. “The major onion growing states are Maharashtra, Madhya Pradesh, Karnataka, Rajasthan, Bihar and Odisha. The Karnataka state stands second position in area (2.49 lakh hectare) and third position in production (20.49 lakh tonnes). The major onion growing districts in the state are Vijayapura, Chitra Durga, Bagalkot, Gadag, Ballary, Dharwad and Haveri” [4].

“Variability in genotypes for yield and its component parameter is the primary factor to consider during selection. Any effective hybridization programme for varietal improvement is primarily dependent on the selection of parents with high variability, so that the desired character combination can be selected to improve quality and also increasing yield production. Furthermore, understanding heredity is critical for selection-based

improvement as it indicates the transmissibility of a character to subsequent generations” [5]. The use of genetically divergent parents in crop improvement is expected to give desirable and superior segregates. This could be achieved by crop improvement.

2. MATERIALS AND METHODS

The present investigation on “Genetic variability, heritability and genetic advance in white onion (*Allium cepa* L.) for growth, bulb yield and its component characters” was conducted at Main Agricultural Research Station, University of Agricultural Sciences, Dharwad during latekharif 2023-24. “Twenty-six genotypes were collected from different institutions and geographical diverse locations and evaluated using randomized complete block design (RCBD) consisting three replications. Five plants were selected randomly from each replication and data were recorded for the characters viz., plant height (cm), number of leaves per plant, leaf length (cm), neck diameter (cm), fresh weight of plant (g), dry weight of plant (g), dry matter content of plant (%), equatorial diameter (cm), polar diameter (cm), bulb index, ten bulb weight (g), average weight of bulb (g), total yield (kg/plot), total yield (t/ha), marketable yield (t/ha), harvest index (%), number of rings per bulb, purple blotch incidence(%), TSS (°Brix), reducing sugar (%), non-reducing sugar (%), total sugar(%) and pyruvic acid (μ moles/ g). Analysis of variance was computed as per the procedures” given by Panse and Sukhatme (1961) and genetic parameters such as mean, range, genotypic and phenotypic characters were analysed as suggested by Burton [6]. Heritability

and genetic advance were worked according to Johanson et al. [7].

3. RESULTS AND DISCUSSION

The nature and degree of genetic variability is one of the most important aspects of any breeding effort. Knowledge of multiple variability characteristics considerably enhances the ability to forecast the degree of variability present in a given set of genetic material, including general mean, Phenotypic Coefficient of Variation (PCV), Genotypic Coefficient of Variation (GCV), range, genetic gain, and broad sense heritability. PCV and GCV estimates were calculated for all of the observed characteristics. "The coefficients of variability are characterized as low, moderate, or high, and they are useful in identifying the level of genetic variability among genotypes and predicting the results of hybridization programs. Heritability is an extremely important parameter for breeders since it indicates how well a genotype may be identified based on its phenotypic expression. Studying genetic improvement with heredity is more important for evaluating the true impacts of selection because heritability alone is insufficient. Genetic factors such as heritability, genotypic coefficient of variation, phenotypic coefficient of variation, and genetic advance (GA) as a percentage of mean were estimated for various onion genotype traits" [7].

The data presented in Table 1 indicates the existence of wide range of phenotypic variability in experimental material. The estimates of PCV were higher than corresponding GCV for all characters studied which indicated that the apparent variation is not only due to genotypes but also due to the influence of environment. The higher estimates of GCV and PCV were recorded for marketable yield (t/ha) (GCV=36.15 %; PCV=37.02 %). Moderate estimates of GCV and higher estimates PCV was registered for Pyruvic acid (μ moles/g) (GCV=19.25 %; PCV=20.80 %). Moderate estimates of GCV and PCV was registered for Polar diameter (cm) (GCV=18.47 %; PCV=18.86 %). Moderate estimate of PCV and lower estimate of GCV was found in the traits Leaf length (cm) at 30 DAT (GCV=9.96 %; PCV=12.08 %), Lowest GCV and PCV were recorded for number of rings per bulb (GCV=5.56 %; PCV=7.94 %).

The higher estimates of GCV and PCV for above characters indicates the existence of sufficient variability among the genotypes for these

characters. Thus, simple selection could be helpful for further improvement. The similar outcome of results was recorded by Gurjar and Singhanian [8], Chattoo et al.[9], Priyanka et al.[10], Amir et al.[11] and Hulagannavar et al.[5].

As the genotypic coefficient of variation does not furnish complete details to determine the variations that are heritable nature. Hence, the heritability estimate become inevitable as this governs the variations due to genetic factors. This heritability helps for direct selection of characters, one should concentrate on these traits which are possessing high heritability. In addition to this, Burton (1952) proposed that the GCV along with heritability estimates will provide the clearer image of the improvement to be expected by phenotypic selection. "The heritability in broad sense comprises both additive and non-additive gene effects" [12].

The estimates of heritability (broad sense) ranges from 48.97 to 95.90 % for various characters under investigation (Table 1). Heritability was maximum for polar diameter (cm) (95.90 %) followed by marketable yield (t/ha) (95.32 %), purple blotch incidence (%) (94.06 %), total yield (t/ha) (93.60 %), average weight of bulb (g) (92.09 %), fresh weight of plant (g) (89.52 %), dry weight of plant (g) (89.08 %), equatorial diameter (cm) (86.64 %), pyruvic acid (μ moles/g) (85.64 %), neck diameter of bulb (cm) (80.93 %), reducing sugar (%) (73.66 %), dry matter content of plant (%) (68.63 %), total sugar (%) (65.47 %), TSS ($^{\circ}$ Brix), (64.47 %), harvest index (%) (63.48 %), number of leaves per plant at harvest(cm) (63.15 %), leaf length at harvest(cm) (59.80 %), bulb index (59.09 %) and plant height at harvest(cm) (52.22 %). While, low estimates of heritability had found for number of rings per bulb (48.97 %). "In this regard earlier studies also reported significant heritability for various parameter among onion genotypes" [13-16].

Genetic advancement is described as an annual rise in performance of a certain characteristic or an annual intensification of that particular trait's performance. The highest genetic advance was recorded for fresh weight of plant (g) (37.47 %) and average weight of bulb (g) (35.17 %). While, moderate values of genetic advance were recorded for purple blotch incidence (%) (15.64 %), marketable yield (t/ha) (12.40 %) and total yield (t/ha) (11.93 %) and Low estimates of genetic advance was recorded for harvest (%)

Table 1. Mean coefficient of variation, heritability (broad sense), genetic advance and genetic advance as per cent of mean of various character of white onion genotypes

Characters	Mean	Range		Variance		Co-efficient of variance		Heritability (Per cent)	Genetic advance (%)	Genetic advance as per cent of mean (Genetic Gain) (%)
		Min.	Max.	Genotypic	Phenotypic	Genotypic	Phenotypic			
X ₁	59.22	44.88	67.68	12.31	23.57	5.92	8.19	52.22	5.22	8.82
X ₂	10.62	7.91	12.18	0.46	0.73	6.40	8.05	63.15	1.11	10.48
X ₃	53.13	40.18	63.47	14.24	24.04	7.10	9.22	59.80	5.98	11.26
X ₄	1.15	0.75	1.43	0.01	0.02	11.50	12.79	80.93	0.24	21.32
X ₅	85.81	47.61	127.09	369.68	412.94	22.40	23.68	89.52	37.47	43.67
X ₆	11.62	6.74	17.92	7.03	7.89	22.81	24.17	89.08	5.15	44.36
X ₇	13.60	11.91	15.87	0.64	0.93	5.89	7.11	68.63	1.36	10.06
X ₈	5.68	3.05	7.44	0.93	1.07	17.03	18.30	86.64	1.85	32.66
X ₉	4.63	2.12	5.97	0.73	0.76	18.47	18.86	95.90	1.72	37.26
X ₁₀	0.81	0.67	0.98	0.002	0.004	6.26	8.15	59.09	0.08	9.92
X ₁₁	65.25	29.55	103.25	316.54	343.73	27.26	28.41	92.09	35.17	53.89
X ₁₂	19.20	6.72	32.25	35.87	38.33	31.18	32.23	93.60	11.93	62.15
X ₁₃	17.06	4.05	28.99	38.02	39.89	36.15	37.02	95.32	12.40	72.70
X ₁₄	75.22	61.50	87.26	18.30	28.83	6.68	7.13	63.48	7.02	9.33
X ₁₅	7.49	6.03	9.07	0.17	0.35	5.56	7.94	48.97	0.60	8.01
X ₁₆	13.49	8.78	17.59	2.14	3.32	10.84	13.51	64.47	2.42	17.94
X ₁₇	2.82	2.05	4.09	0.15	0.20	13.91	16.21	73.66	0.69	24.60
X ₁₈	5.96	4.35	8.15	0.42	0.64	10.86	13.43	65.47	1.08	18.11
X ₁₉	3.90	2.27	5.32	0.56	0.65	19.25	20.80	85.64	1.43	36.70
X ₂₀	23.24	9.48	41.92	61.36	65.23	33.70	34.75	94.06	15.64	67.34

Note: X₁-plant height (cm) at harvest, X₂- no. of leaves per plant at harvest, X₃-leaf length (cm) at harvest, X₄-neck diameter of bulb (cm), X₅-fresh weight of plant (g), X₆-dry weight of plant (g), X₇- dry matter content of plant (%), X₈-equatorial diameter (cm), X₉-polar diameter (cm), X₁₀-bulb index, X₁₁- average weight of bulb (g), X₁₂-total yield (t/ha), X₁₃-marketable yield (t/ha), X₁₄-harvest index (%), X₁₅-number of rings per bulb, X₁₆- tss (°brix), X₁₇-reducing sugar (%), X₁₈-total sugar (%), X₁₉-pyruvic acid (μ moles/g), X₂₀- purple blotch incidence (%)

index (7.02 %), leaf length at harvest (cm) (5.98 %), dry weight of plant (g) (5.15 %), plant height at harvest(cm) (5.22 %), TSS (°Brix) (2.42 %), equatorial diameter (cm) (1.85 %), polar diameter (cm) (1.72 %), pyruvic acid (μ moles/g) (1.43 %), dry matter content of plant (%) (1.36 %), number of leaves per plant at harvest (cm) (1.11 %), total sugar (%) (1.08 %), reducing sugar (%) (0.69 %), number of rings per bulb (0.60 %), neck diameter of bulb (cm) (0.24 %), and bulb index (0.08 %). Similarly, Dhotre et al. [13], Hosamani et al. [17] and Hulagannavar et al.[5] also showed variations of genetic advance for various parameter among the onion genotypes.

The data elicited on genetic advance as per cent of mean(genetic gain) showed maximum value for marketable yield (t/ha) (72.70 %), purple blotch incidence (%) (67.34 %), total yield (t/ha) (62.15 %), average weight of bulb (g) (53.89 %), dry weight of plant (g) (44.36 %) and fresh weight of plant (g) (43.67 %). While, moderate values of genetic advance as per cent of mean was noticed for polar diameter (cm) (37.26 %) followed by pyruvic acid (μ moles/g) (36.70 %), equatorial diameter (cm) (32.66 %), reducing sugar (%) (24.60 %), neck diameter of bulb (cm) (21.32 %), total sugar (%) (18.11 %), TSS (°Brix) (17.94 %), leaf length at harvest (cm) (11.26 %), number of leaves per plant at harvest(cm) (10.48 %) and dry matter content of plant (%) (10.06 %) and lowest values of genetic advance as per cent of mean was registered for harvest index (%) (9.33 %), bulb followed by index (9.92 %), plant height at harvest (cm) (8.82 %), and number of rings per bulb (8.01 %)(Table 1). In parallel to our investigation, earlier studies also reported the significant difference in genetic advance as per cent of mean among the different onion cultivars for various parameters (Manjunathagowda et al., [16] Singh et al., [18], and Hulagannavar et al., [5,19.]

4. CONCLUSION

For every characteristic under investigation, significant genotypic and phenotypic differences were found. A high genotypic coefficient of variation was found for yield, suggesting that there is a good deal of variability and that the environment has little effect on how it manifests itself. For the variables marketable yield (t/ha) and total yield (t/ha), the aforementioned genetic parameters—high heritability estimates combined with genetic advance—were registered based on the current experiment. It reveals the presence of additive gene action in expression of

these traits and are more definitive for potential direct selection these traits.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous, Area, production and productivity of onion; 2020. Available:www.faostat.org
2. Manjunath U, Hiremath SM. Genetic variability, heritability and genetic advance in onion (*Allium cepa* L.) for bulb yield and its component characters. Pharma Innov. 2022;11(11):2570-2.
3. Anonymous, Area, Production and Productivity of Onion; 2021. Available: www. nhrdf .com.
4. Anonymous, Area, production and productivity of onion; 2022. Available:https://des.karnataka.gov.in
5. Hulagannavar P, Patil B, Gunnaiah R and Cholin S. Estimates of variability, heritability, genetic advance for yield and its quality traits in onion (*Allium cepa* L.) genotypes. International Journal of Environment and Climate Change. 2023;13(10):1758-1770.
6. Burton GW, Devane EH. Estimating heritability in tall fescue (*Festuca arundinacea*) from replicated clonal material. Agronomy Journal. 1953;45:478-481.
7. Johnson HW, Robinson HF, Comstock RF. Estimates of genetic and environmental variability of soybeans. Agronomy Journal. 1955;47(1):317-318.
8. Gurjar RS, Singhania DL. Genetic variability, correlation and path analysis of yield components in onion. Indian Journal of Horticulture. 2006;63(1):53-58.
9. Chattoo MA, Angrej A. Genetic variability, interrelationship and path analysis for yield and yield related traits in onion (*Allium cepa* L.) under temperate condition in Kashmir valley. Plant Archives. 2015;15(2):1161-1165.

10. Priyanka A, Dod VN, Sharma M. Variability studies in *rabi* onion (*Allium cepa* L) for yield and yield contributing traits. International Journal of Farm Science. 2017;7(1):123-126.
11. Amir A, Sharangi AB, Bal S, Upadhyay TK, Khan MS, Ahmad I, Alabdallah NM, Saeed M, Thapa U. Genetic variability and diversity in red onion (*Allium cepa* L.) Genotypes: Elucidating morpho-horticultural and quality perspectives, Horticulturae. 2023;9(9):1-17.
12. Hanson WD, Robinson HF, Comstock RE. Biometrical studies of yield in segregating population of Korean lespedeza. Agronomy Journal. 1956;48(6):268-278.
13. Dhotre M, Allolli TB, Athani SI, Halemani LC. Genetic variability, character association and path analysis studies in *kharif* onion (*Allium cepa* L.). Asian Journal of Horticulture. 2017;5(1):143-146.
14. Sharma PK, Singh A, Duhan DS, Kishor N, Barar NS. Genetic variability, heritability and genetic advance in onion (*Allium cepa* var. *cepa* L.). International Journal of Pure and Applied Bioscience. 2017;5(6):740-743.
15. Srivastav G, Vikram B, Prasad VM. Studies on multiple correlation between bulb yield, growth and yield attributes in different genotypes of onion (*Allium cepa* L.) under Allahabad agro-climatic condition. Journal Pharmacognosy and Phytochemistry. 2017; 6(6): 793-798.
16. Manjunathagowda DC, Anjanappa M, Jayaswall K, Venugopalan R, Kumar A, Shankarappa KS, Lingaiah HB. Perspective and application of molecular markers linked to the cytoplasm types and male-fertility restorer locus in onion (*Allium cepa*). Plant Breeding. 2021;140(5):732-44.
17. Hosmani RM, Patil BC, Ajjappalavara. Genetic variability and character association studies in onion (*Allium cepa*L.). Karnataka Journal of Agricultural Sciences. 2010;23(2):302- 305.
18. Singh D, Trivedi J, Sharma PK, Lodhi Y. Evaluation of different onion (*Allium cepa*L.) genotypes for growth, yield and quality parameters under Chhattisgarh plain region. The Pharma Innovation Journal. 2021;10(9):1646-1650.
19. Panse VG, Sukhatme PV. Statistical methods for agricultural workers (2ndEdn.). ICAR Publication, New Delhi, India; 1961.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/121982>