



Comparison between Pharmaco-invasive Strategy and Primary Percutaneous Coronary Intervention According to Percutaneous Coronary Intervention Related Delay

**Khaled Nasr Akl^{1*}, Medhat Mohamed Ashmawy¹, Magdy Mohamed Elmasry¹,
Hanan Kamel Kassem¹ and Ayman Mohamed Elsaied¹**

¹*Department of Cardiovascular Medicine, Faculty of Medicine, Tanta University, Egypt.*

Authors' contributions

This work was carried out in collaboration among all authors. Author KNA designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors MMA and MME managed the analyses of the study. Authors HKK and AME managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMMR/2020/v32i730455

Editor(s):

(1) Prof. Franciszek Burdan, Medical University of Lublin, Poland.

Reviewers:

(1) Maruyama Kyushu, Japan.

(2) Akinlade Olawale Mathias, Nigeria.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/57168>

Original Research Article

Received 18 March 2020

Accepted 24 May 2020

Published 30 May 2020

ABSTRACT

Aim: This study aimed to compare between the effect of pharmacoinvasive strategy (PI) & primary percutaneous coronary intervention (P-PCI) according to PCI related delay (door to balloon) on the mortality and morbidity during in-hospital stay and after 30-day follow up. Moreover, left ventricular systolic function was assessed by two-dimensional echocardiography at cardiology department, Tanta University.

Patients and Methods: The study was conducted on 300 patients that were divided into 2 main groups. Group A consisted of patients had primary PCI as reperfusion therapy and further divided into three groups according to PCI related delay (door to balloon). Group A1, PCI-related delay is ≤ 60 minute (92 patients). Group A2, PCI-related delay is >60 to ≤ 90 minute. (54 patients). Group A3, PCI-related delay is >90 minute (78 patients). The second group (group B), include patients who under pharmacoinvasive strategy, PCI within 24 hour after thrombolysis (76 patients). In

*Corresponding author: E-mail: khaledakl510@gmail.com;

the second group, coronary angiography was done immediately in cases of failed thrombolysis and for successful thrombolysis; coronary angiography was performed within 3 – 24 hours.

Results: During hospital stay, more patients in group A3 died than those of group B or group A1, A2 with no statistical significance. In addition, more patients in group A3 showed heart failure symptoms with statistical significance than those of group B, A1 and A2. Bleeding complications occurred significantly more in group B. During follow up visits more patients in group A3 complained of heart failure symptoms with statistical significance than those of group B, A1, A2 patients.

Conclusion: Primary PCI without door to balloon time delay (≤ 90 minutes) was encouraged and had the best results on morbidity and mortality. Also, pharmacoinvasive strategy was encouraged as being better than primary PCI when door to balloon time showed marked delay (< 90 minutes).

Keywords: Pharmaco-invasive strategy; primary PCI.

1. INTRODUCTION

ST elevation myocardial infarction (STEMI) is a clinical syndrome defined by characteristic symptoms of myocardial ischemia accompanied with persistent electrocardiographic ST elevation and the subsequent release of biomarkers due to myocardial necrosis. The most common triggering event is the disruption of an atherosclerotic plaque in an epicardial coronary artery, which leads to a clotting cascade, resulting in total obstruction of the coronary artery. The current definitive treatment for STEMI is reperfusion therapy consisting of percutaneous coronary intervention and thrombolysis, as recommended by the American College of Cardiology Foundation/American Heart Association and the European Society of Cardiology [1,2].

STEMI is often caused by complete obstruction of an artery by a blood clot (thrombus). As soon as the coronary blood supply is interrupted, heart muscle (Myocardium) begins to be damaged, and the longer the blood supply is obstructed the greater the myocardial damage. In animal models nearly half of potentially viable myocardium is lost within 1 hour, and two-thirds lost within 3 hours, of experimental coronary artery occlusion [3].

The objectives of treatment are to restore coronary blood supply flow (reperfusion) as soon as possible after the onset of symptoms of acute STEMI. Reperfusion can be achieved by mechanical techniques (coronary angioplasty, thrombus extraction catheters, stenting) that are grouped under the overarching term "Primary percutaneous coronary intervention" (P-PCI), or by the use of fibrinolytic drugs that lyse the coronary thrombus [3,4].

Primary percutaneous coronary intervention (P-PCI) is the recommended reperfusion therapy rather than fibrinolysis, if performed within 90-

120 minute from the first medical contact in an expert 24/7 facility. If PCI cannot be performed within 90 to 120 minute, then thrombolysis, preferably within 30 minutes of arrival at the hospital, is recommended. Fibrinolytic therapy is recommended in patients without contraindications. The original studies supporting this view however compared primary-PCI with in-hospital fibrinolysis only [5,6].

An important subsequent development has been pharmacoinvasive (PI) strategy (early fibrinolysis with rescue PCI if fibrinolytic fail and with subsequent early angiography/PCI following fibrinolytic success). Some data support pharmacoinvasive strategy as being equal or better than primary PCI especially when door to balloon time show marked delay [7,8].

Other studies indicate that there is a direct relation between infarct size and that mortality rates increase the longer it takes to deliver primary PCI. By inference any advantage of primary PCI over fibrinolysis may become attenuated the longer the PCI related delay. Thus, pharmacoinvasive strategy can provide an alternative strategy to primary PCI [9,10].

2. PATIENTS AND METHODS

2.1 Patients

This study was conducted at the department of cardiovascular medicine, Tanta University hospital, it was carried out on 300 patients diagnosed with acute STEMI. Patients were divided into two main groups according to the method of reperfusion they received.

Group A: included patients who underwent primary percutaneous coronary intervention strategy (224 patients). Then, they were divided into three categories according to PCI-related delay (Door to balloon).

Group A1: PCI-related delay (Door to balloon) was ≤ 60 minute. (92 patients). **Group A2:** PCI-related delay (Door to balloon) was >60 to ≤ 90 minute. (54 patients). **Group A3:** PCI-related delay (Door to balloon) was > 90 minute. (78 patients)

Group B: included patients who underwent PCI within 24 hour after thrombolysis (pharmacoinvasive strategy). (76 patients)

Inclusion criteria: Patients presented within 12 hours of symptom onset with STEMI and treated with primary PCI or pharmacoinvasive strategy.

Exclusion criteria:

- Patients with myocardial infarction (MI) refused the previous two methods of intervention.
- Patients with MI and late presentation (more than 12 hours of symptom onset).
- Patients with MI with non-obstructive coronary artery disease (MINOCA) as acute myocarditis excluded by coronary angiography.

2.2 Methods

All patients were subjected to the following:

- 1- An informed consent was taken from all participants.
- 2- **History taking:** including personal history, presence of risk factors, past cardiac history and history of any comorbidities.
- 3- **Clinical examination:** including vital signs, General examination on the abdomen, chest, head, neck, and both the upper and lower limbs and local cardiac examination.
- 4- 12 leads surface electrocardiography (ECG).
- 5- **Blood sampling:** Serum cardiac biomarkers, complete blood count, random blood sugar serum, urea & creatinine, lipid profile.
- 6- Reperfusion either through primary percutaneous intervention for Infarct related artery or through pharmacoinvasive technique:

In pharmacoinvasive technique, Percutaneous coronary intervention was performed either immediately after failure of thrombolytic therapy (rescue PCI) or within 3 -24 hours after criteria of successful thrombolysis (routine early angiography/PCI strategy).

7- **Echocardiography:** full two-dimensional (2D) and M mode echocardiographic study in the standard views (parasternal long axis, short axis, apical four and apical two chambers) was done to assess left ventricular (LV) systolic function, resting segmental wall motion abnormalities (RSWMA) and any mechanical complications.

8- The study compared between those groups in the acute stage during hospitalization of the patients according to the following: a- Clinical outcomes: major adverse cardiac events (MACE) as mortality, heart failure, re-infarction. Also, bleeding complication, neurological complication and contrast induced nephropathy. b- Angiographic findings (base line and final TIMI score and angiographic complications as dissection and perforation). c- LV systolic function assessment by echocardiography.

9- Short-term follow up after one month for: a- Clinical outcomes: major adverse cardiac events (MACE) as mortality, heart failure, re-infarction and cerebrovascular accidents. b- LV systolic function assessment by echocardiography.

2.3 Statistical Analysis of the Data

The analysis was calculated by SPSS version 25. The qualitative parameters were described by number of frequency and percentage while the quantitative variables were described by mean, standard deviation and range. In addition, the comparison of independent quantitative variables was calculated by Anova with Tukey test in post hoc analysis. However, comparison between two qualitative variables was done by Chi square, Fisher's exact fisher and Monte Carlo tests.

3. RESULTS

3.1 Comparison between the Study Groups According to the Demographic Data, Risk Factors and Clinical Presentation

Males represented 73.3% of patients, while females represented 26.6% of patient presented by STEMI with a ratio of 2.75:1. The age of the study population ranged from 28-88 years. In this

study 103 patients were diabetics, and 129 were hypertensive, while 138 were active smokers and 46 were addicts. In this study 186 of the study population presented by anterior STEMI in which left anterior descending (LAD) was the culprit lesion, 98 patients presented by inferior STEMI and 10 patients presented by lateral STEMI, 6 patients presented by isolated posterior STEMI. Also 224 patient in our study presented by Killip class I while 39 patients presented by Killip class II and 36 patients presented by Killip Class III, IV.

3.2 Comparison between the Study Groups according to In-Hospital Adverse Effects

Regarding in-hospital mortality: There was no significant differences between four groups.

Regarding congestive heart failure symptoms: There was statistically significant difference between the four groups with marked incidence of congestive heart failure (CHF) in group A3.

Bleeding complication showed statistically significant difference between the four groups with marked incidence in group B.

3.3 Comparison between the Study Groups According to Follow-Up Adverse Effects (30 Days)

During follow up visit, there were no statistically significant difference between groups regarding all-cause mortality.

But, there was statistically significant difference between the four groups regarding congestive heart failure with marked incidence of heart failure in group A3.

For the median ejection fraction during admission there was statistically significant difference between the four groups with marked decrease in systolic function in patients in group A3.

For assessment of left ventricular systolic function after one month there was statistically significant difference between the four groups with marked decrease in systolic function in patients in group A3.

Table 1. Comparison between the studied groups according to clinical presentation

		Group A1 N (%) N=92	Group A2 N (%) N=54	Group A3 N (%) N=78	Group B N (%) N=76	X ²	P Value
Killip	1	78 (84.8%)	38 (70%)	52 (66.7%)	57 (75%)	9.091	0.168
	2	7 (7.6%)	9 (16.7%)	12 (15.4%)	11 (14.5%)		
	3 and 4	7 (7.6%)	7 (13%)	14 (17.9%)	8 (10.5%)		
Location of MI	Anterior	63 (68.47%)	25 (46.2%)	48 (61.53%)	50 (65.78%)	11.625	0.226
	Inferior	23 (25%)	26 (48.1%)	27 (34.61%)	22 (28.94%)		
	Lateral	3 (3.261%)	2 (3.704%)	3 (3.846%)	2 (2.631%)		
	Isolated posterior	3 (3.261%)	1 (1.852%)	0 (0%)	2 (2.631%)		

Table 2. Comparison between the study groups according to in-hospital adverse effects

		Group A1 N (%) N=92	Group A2 N (%) N=54	Group A3 N (%) N=78	Group B N (%) N=76	X ²	P Value
Mortality		4 (4.348%)	5 (9.259%)	10 (12.8%)	6 (7.895%)	4.049	0.256
Bleeding	Minor bleed	7 (7.6%)	7 (13%)	8 (10.3%)	17 (22.4%)	12.95	0.043*
	Major bleed	2 (2.2%)	1 (1.9%)	2 (2.6%)	5 (6.6%)		
Heart failure		10 (10.9%)	10 (18.5%)	23 (29.5%)	12 (15.8%)	10.23	0.017*
Cardiogenic shock		4 (4.348%)	4 (7.407%)	11 (14.1%)	5 (6.579%)	5.848	0.119

Table 3. Comparison between the study groups according to follow-up adverse effects (30 days)

	Group A1 N (%) N=82	Group A2 N (%) N=44	Group A3 N (%) N= 58	Group B N (%) N=61	X ²	P Value
Follow up Mortality	2 (2.4%)	3 (6.8%)	5 (8.6%)	4 (6.6%)	2.72	0.5
Follow up HF	6 (7.3%)	5 (11.4%)	13 (22.4%)	5 (8.2%)	8.6	0.035*

Table 4. Comparison between the studied groups according to EF assessment by Echocardiography before discharge&during follow up and RSWMA and Dilated dimensions

	Group A1 mean±SD (range) N=92	Group A2 mean±SD (range) N=54	Group A3 mean±SD (range) N=78	Group B mean±SD (range) N=76	F	P Value	Post Hoc Test
ECHO (EF) %	46.48 ± 8.69 (30-66)	45.57 ± 9.9 (20-64)	40.52 ± 9.2 (20-68)	43.8 ± 9.5 (28-65)	6.844	<0. 001*	P1 0. 942 P2<0. 001* P3 0. 135 P4 0. 01* P5 0. 633 P6 0. 04*
Follow up EF%	50.92 ± 7.87 (35-66)	50.32 ± 10.589 (20-65)	44.4 ± 9.62 (18-68)	49.79 ± 9.1 (30-65)	6.57	<0. 001*	P1 0. 985 P2<0. 001* P3 0. 88 P4 0. 007* P5 0. 991 P6 0. 007*
	Group A1 N (%) N= 92	Group A2 N (%) N=54	Group A3 N (%) N=78	Group B N (%) N= 76	X ²		P Value
ECHO (RWMA)	87(87%)	49(90.7%)	71(91%)	66(86.8%)	1.183		0.75
ECHO (Dilated)	16(17.4%)	15(28.8%)	29(37. 2%)	11(14.5%)	13.95		0.003*

4. DISCUSSION

Males represented 73.3% of patients, while females represented 26.6% of patient presented by STEMI with a ratio of 2.75:1. The age of the study population ranged from 28-88 years.

A study conducted by *STREAM TRIA* [11] in which the ratio of males to females having MI was 4:1 in the study population. Also, this came in agreement with the American heart association statistical annual updated report by Mozaffarian et al. [12] that found that STEMI is more prevalent in men than women.

In this study 186 of the study population presented by anterior STEMI in which LAD was the culprit lesion (62%), 98 patients presented by inferior STEMI (32.6%) and 10 patients presented by lateral STEMI (3.3%),6 patients presented by isolated posterior STEMI (2%).Also 224 patient in our study presented by Killip class I (74.6%) while 39 patients (13%) presented by Killip class II and 36 patients (12%) presented by Killip Class III,IV.

This came in agreement by *STREAM trial* [11] in which the majority of cases presented by anterior STEMI and patients presenting by Killip class I represented then majority of their study population 94% of all patients.In the study conducted by Gershlick et al. [13] anterior STEMI represented about 47% of MI and Killip class I

represented the majority of their study population.

Study groups were compared regarding base line TIMI flow in coronary angiography. In group B, treated with fibrinolytic agents (30.26%) of patients achieved TIMI 0,I flow. While (69.73%) of patients achieved either TIMI flow II, III. This result was due to use of thrombolytic therapy before undergoing coronary angiography.But as would be expected in group A,the majority of cases were TIMI 0 with percentage of 69.5%,70.3%,73,0% of cases in groups A1,A2,A3 respectively.After PCI, patency rates were high in all study groups with final TIMI III achieved in (91.3%) in group A1,(79.6%) in group A2,(70.5%) in group A3, (88.1%)in group B. With marked incidence of no-reflow in group A3 (7.6%) of cases.

In *STREAM trial*, [11] in the group treated by pharmacoinvasive technique most patients presented by base line TIMI III (58.5%)of cases, while in the group treated by primary PCI most patients achieved base line TIMI 0 (59.3%). But the final TIMI III flow was achieved similarly in the group treated by pharmacoinvasive technique and group treated by primary PCI 91% and 92% respectively. Also the study conducted by Gershlick et al. [13] initial TIMI III flow for group treated by primary PCI is 21.4% of patients. And 58.6% of patients treated by pharmacoinvasive strategy.While the final TIMI III

flow was 92.3% in group treated by primary PCI and 91.6% in patients treated by pharmacoinvasive strategy.

Also in the *FAST-MI trial* [14] initial TIMI III flow for group treated by primary PCI is 18% of patients. And 37% of patients treated by pharmacoinvasive technique. While the final TIMI III flow was 89% in group treated by primary PCI and 84% in patients treated by pharmacoinvasive strategy. Regarding in-hospital mortality: In group A1, 4 patients died during hospital stay (4.3%). In group A2, 5 patients (9.2%). In group A3, 10 patients (12.8%). And in group B 6 patients (7.8%) there were no significant differences between groups.

Congestive heart failure symptoms occurred in group A1, 10 patients (10.8%). In group A2, 10 patients (18.518%). In group A3, 21 patients (26.923%). And in group B, 12 patients (15.7%). There was statistically significant difference between the four groups with marked incidence of CHF in group A3.

Bleeding complication occurred more in the pharmacoinvasive arm compared with primary PCI arm, 17 patients (22.3%) had minimal & minor bleeding and 5 patients (6.5%) had major bleeding within group B. There was statistically significant difference between the four groups with marked incidence in pharmacoinvasive strategy.

During follow up visit, there were no marked differences between groups regarding all-cause mortality. In group A1, 2 patients died (2.4%). In group A2, 3 patients (6.8%). In group A3, 4 patients (6.8%). And in group B, 4 patients (6.5%).

But, there were marked differences between groups regarding congestive heart failure. In group A1, 6 patients (7.3%). In group A2, 5 patients (11.3%). In group A3, 11 patients (18.9%). And in group B, 5 patients (8.1%). There was statistically significant difference between the four groups due to marked incidence of heart failure in group A3.

In the study conducted by Gershlick et al. [13] compared outcome in patients treated with pharmacoinvasive therapy (PI) with primary PCI (P-PCI) according to PCI related delay (P-RD) and categorized patients in three groups, first with PCI related delay ≤ 55 minute, second with PCI related delay $> 55-97$ minute and third group with PCI related delay > 97 minute and compared them with those undergoing pharmacoinvasive strategy.

Comparing death, congestive heart failure, cardiogenic shock and myocardial infarction in Pharmacoinvasive therapy with primary PCI arms occurred in 10.6% versus 10.3% (≤ 55 minute); 13.9% versus 17.9% ($> 55-97$ minute) and 13.5% versus 16.2% (> 97 minute), respectively. For P-RD ≤ 55 min, fewer events tended to occur with P-PCI than PI [13].

Conversely, as PCI-related delay (P-RD) increased to > 55 minute, patients with Pharmacoinvasive therapy had better outcomes than primary PCI (P-PCI), suggesting advantages with PI when P-RD delay [13].

This came in agreement with *STREAM trial*, [11] which compared outcomes in patients treated with pharmacoinvasive therapy or Primary PCI presenting within 3 hours after symptom onset, unable to undergo P-PCI within 1 hour. The primary end point was a composite of death, shock, congestive heart failure, or reinfarction up to 30 days. The primary end point occurred in (12.4%) in the Pharmacoinvasive group and in (14.3%) in the primary PCI group. More intracranial hemorrhages occurred in the Pharmacoinvasive group than in the primary PCI group.

In the *FAST-MI trial*, [14] they assessed 5-year mortality in STEMI patients from the *French registry of Acute ST-elevation or non-ST elevation Myocardial Infarction* (FAST-MI) 2005 according to use and type of reperfusion therapy. Of 1492 STEMI patients with first call < 12 hours from onset, 447 (30%) received fibrinolysis (66% pre-hospital; 97% with subsequent angiography, 84% with subsequent PCI), 583 (39%) had primary PCI and 462 (31%) received no reperfusion. There was a numerical excess of reinfarction, stroke, and ventricular fibrillation with the fibrinolytic-based strategy, and an excess of cardiogenic shock with primary PCI. However, none of the in-hospital complications differed significantly for the two reperfusion strategies. In the FAST-MI trial major bleeding complication occurred more with the primary PCI arm with no statistical difference. While in 5-year follow up, Five-year survival was high in patients who had received reperfusion therapy with either primary PCI, or a pharmaco-invasive approach, with approximately two-thirds of the patients receiving fibrinolytic treatment in the pre-hospital setting. As expected, patients who did not get reperfusion therapy had a much higher mortality. When comparing the two reperfusion strategies, the results achieved with the pharmaco-invasive approach were at least as good as those with an

intended primary PCI strategy. Also, the *Comparison of Angioplasty and Prehospital Thrombolysis in Acute Myocardial Infarction (CAPTIM) trial*, [7] has told that prehospital fibrinolytic therapy with the patients brought to PCI-capable centers and with one third undergoing rescue angioplasty, could do at least as well as primary PCI up to 5 years after the initial episode.

5. CONCLUSION

Primary PCI without door to balloon time delay (≤ 90 minutes) was encouraged and had the best results on morbidity and mortality. But in our daily clinical practice pharmacoinvasive strategy was considered safe alternative to primary PCI. Especially, considering logistical issues and delay in the initiation of management and the results of our study supported pharmacoinvasive strategy as being better than primary PCI when door to balloon time showed marked delay (> 90 minutes).

6. LIMITATIONS OF THE STUDY

The study had some potential limitations such as; small size of study population, which was due to many factors, one of them that not all patients were willing to the idea of follow up after one month, also a lot of cases came with late presentation after the accepted window of thrombolytic therapy, others refused doing PCI at our center due to logistic or cultural issues.

In addition, many patients who received thrombolytic therapy with signs of successful reperfusion underwent coronary angiography later on after discharge due to financial reasons.

Another limitation was the short period assigned for follow up which didn't allow the appearance of results for mortality, reinfarction & re-hospitalization. The chosen period was one month only to prevent fallacies in the results because mostly after one month the patients underwent elective PCI for other coronary lesions so this may affect the results.

Also the use of Simpson's method, M-mode might not be of the same accuracy in assessment the global & regional LV systolic function as the newest techniques such as speckle tracking & strain and strain rate.

CONSENT

An informed consent was taken from all participants.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. O'Gara PT, Kushner FG, Ascheim DD, Casey DE, Chung MK, de Lemos JA. ACCF/AHA guideline for the management of ST-elevation myocardial infarction: A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: Developed in collaboration with the American College of Emergency Physicians and Society for Cardiovascular Angiography and Interventions. *Catheterization and cardiovascular interventions: Official journal of the Society for Cardiac Angiography & Interventions*. 2013;82(1):E1-27.
2. Steg PG, James S. ESC Guidelines on acute myocardial infarction (STEMI). *European Heart Journal*. 2012;33:2501-2.
3. Antonucci D, Valenti R, Migliorini A, Moschi G, Trapani M, Buonamici P. Relation of time to treatment and mortality in patients with acute myocardial infarction undergoing primary coronary angioplasty. *The American Journal of Cardiology*. 2002; 89(11):1248-52.
4. Brodie BR, Hansen C, Stuckey TD, Richter S, Versteeg DS, Gupta N. Door-to-balloon time with primary percutaneous coronary intervention for acute myocardial infarction impacts late cardiac mortality in high-risk patients and patients presenting early after the onset of symptoms. *Journal of the American College of Cardiology*. 2006; 47(2):289-95.
5. Fernandez-Aviles F, Alonso JJ, Castro-Beiras A, Vazquez N, Blanco J, Alonso-Briales J. Routine invasive strategy within 24 hours of thrombolysis versus ischaemia-guided conservative approach for acute myocardial infarction with ST-segment elevation (GRACIA-1): A randomised controlled trial. *Lancet* (London, England). 2004;364(9439):1045-53.
6. Keeley EC, Boura JA, Grines CL. Primary angioplasty versus intravenous thrombolytic

- therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. *Lancet* (London, England). 2003; 361(9351):13-20.
7. Bonnefoy E, Steg PG, Boutitie F, Dubien PY, Lapostolle F, Roncalli J. Comparison of primary angioplasty and pre-hospital fibrinolysis in acute myocardial infarction (CAPTIM) trial: A 5-year follow-up. *European Heart Journal*. 2009;30(13): 1598-606.
 8. Westerhout CM, Bonnefoy E, Welsh RC, Steg PG, Boutitie F, Armstrong PW. The influence of time from symptom onset and reperfusion strategy on 1-year survival in ST-elevation myocardial infarction: A pooled analysis of an early fibrinolytic strategy versus primary percutaneous coronary intervention from CAPTIM and WEST. *American Heart Journal*. 2011;161(2):283-90.
 9. Cannon CP, Gibson CM, Lambrew CT, Shoultz DA, Levy D, French WJ. Relationship of symptom-onset-to-balloon time and door-to-balloon time with mortality in patients undergoing angioplasty for acute myocardial infarction. *JAMA*. 2000; 283(22):2941-7.
 10. Stone GW, Dixon SR, Grines CL, Cox DA, Webb JG, Brodie BR. Predictors of infarct size after primary coronary angioplasty in acute myocardial infarction from pooled analysis from four contemporary trials. *The American Journal of Cardiology*. 2007; 100(9):1370-5.
 11. Armstrong PW, Gershlick AH, Goldstein P, Wilcox R, Danays T, Lambert Y. Fibrinolysis or primary PCI in ST-segment elevation myocardial infarction. *New England Journal of Medicine*. 2013; 368(15):1379-87.
 12. Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M. Heart Disease and Stroke Statistics--2015 Update: A Report From the American Heart Association. *Circulation*. 2014;131:29-322.
 13. Gershlick AH, Westerhout CM, Armstrong PW, Huber K, Halvorsen S, Steg PG, Wilcox RG. Impact of a pharmacoinvasive strategy when delays primary PCI are prolonged. *Heart*. 2015;101(9):692-698.
 14. Danchin N, Puymirat E, Steg PG, Goldstein P, Schiele F, Belle L. Five-Year Survival in Patients With ST-Segment-Elevation Myocardial Infarction According to Modalities of Reperfusion Therapy: The French Registry on Acute ST-Elevation and Non-ST-Elevation Myocardial Infarction (FAST-MI) 2005 Cohort. *Circulation*. 2014;129(16):1629-36.

© 2020 Akl et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/57168>